

146. Hafner J, Schaad I, Schneider E, Seifert B, Burg G, Cassina PC. Leg ulcers in peripheral arterial disease (arterial leg ulcers): impaired wound healing above the threshold of chronic critical limb ischemia. *J Am Acad Dermatol* 2000;43:1001-8.
147. Fontaine R, Kim M, Kieny R. [Surgical treatment of peripheral circulation disorders]. *Helv Chir Acta* 1954;21:499-533.
148. Meggitt B. Surgical management of the diabetic foot. *Br J Hosp Med* 1976;16:227-32.
149. Wagner FW Jr. The dysvascular foot: a system for diagnosis and treatment. *Foot Ankle* 1981;2:64-122.
150. Second European Consensus Document on chronic critical leg ischemia. *Circulation* 1991;84(Suppl):IV1-26.
151. Rutherford RB, Baker JD, Ernst C, Johnston KW, Porter JM, Ahn S, et al. Recommended standards for reports dealing with lower extremity ischemia: revised version. *J Vasc Surg* 1997;26:517-38.
152. Armstrong DG, Lavery LA, Harkless LB. Validation of a diabetic wound classification system. The contribution of depth, infection, and ischemia to risk of amputation. *Diabetes Care* 1998;21:855-9.
153. Macfarlane RM, Jeffcoate WJ. Classification of diabetic foot ulcers: the S(AD) SAD system. *Diabetic Foot* 1999;2:123-6.
154. Dormandy JA, Rutherford RB. Management of peripheral arterial disease (PAD). TASC Working Group. TransAtlantic Inter-Society Consensus (TASC). *J Vasc Surg* 2000;31(Pt 2):S1-296.
155. Schaper NC. Diabetic foot ulcer classification system for research purposes: a progress report on criteria for including patients in research studies. *Diabetes Metab Res Rev* 2004;20(Suppl 1):S90-5.
156. Norgren L, Hiatt WR, Dormandy JA, Nehler MR, Harris KA, Fowkes FG. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *J Vasc Surg* 2007;45(Suppl S):S5-67.
157. Martinez-De Jesus FR. A checklist system to score healing progress of diabetic foot ulcers. *Int J Low Extrem Wounds* 2010;9:74-83.
158. Lipsky BA, Berendt AR, Cornia PB, Pile JC, Peters EJ, Armstrong DG, et al. 2012 Infectious Diseases Society of America clinical practice guideline for the diagnosis and treatment of diabetic foot infections. *Clin Infect Dis* 2012;54:e132-73.
159. Adam DJ, Beard JD, Cleveland T, Bell J, Bradbury AW, Forbes JF, et al. Bypass versus angioplasty in severe ischaemia of the leg (BASIL): multicentre, randomised controlled trial. *Lancet* 2005;366:1925-34.
160. Bradbury AW, Adam DJ, Bell J, Forbes JF, Fowkes FG, Gillespie I, et al. Bypass versus Angioplasty in Severe Ischaemia of the Leg (BASIL) trial: analysis of amputation free and overall survival by treatment received. *J Vasc Surg* 2010;51(Suppl):18S-31S.
161. Shishehbor MH, Hammad TA, Zeller T, Baumgartner I, Scheinert D, Rocha-Singh KJ. An analysis of IN.PACT DEEP randomized trial on the limitations of the societal guidelines-recommended hemodynamic parameters to diagnose critical limb ischemia. *J Vasc Surg* 2016;63:1311-7.
162. Conte MS, Geraghty PJ, Bradbury AW, Hevelone ND, Lipsitz SR, Moneta GL, et al. Suggested objective performance goals and clinical trial design for evaluating catheter-based treatment of critical limb ischemia. *J Vasc Surg* 2009;50:1462-73.e1-3.
163. Beropoulos E, Stavroulakis K, Schwindt A, Stachmann A, Torsello C, Bisdas T. Validation of the Wound, Ischemia, foot Infection (WIFI) classification system in nondiabetic patients treated by endovascular means for critical limb ischemia. *J Vasc Surg* 2016;64:95-103.
164. Darling JD, McCallum JC, Soden PA, Guzman RJ, Wyers MC, Hamdan AD, et al. Predictive ability of the Society for Vascular Surgery Wound, Ischemia, and foot Infection (WIFI) classification system after first-time lower extremity revascularizations. *J Vasc Surg* 2017;65:695-704.
165. Mathioudakis N, Hicks CW, Canner JK, Sherman RL, Hines KF, Lum YW, et al. The Society for Vascular Surgery Wound, Ischemia, and foot Infection (WIFI) classification system predicts wound healing but not major amputation in patients with diabetic foot ulcers treated in a multidisciplinary setting. *J Vasc Surg* 2017;65:1698-705.e1.
166. Ward R, Dunn J, Clavijo L, Shavelle D, Rowe V, Woo K. Outcomes of critical limb ischemia in an urban, safety net hospital population with high WIFI amputation scores. *Ann Vasc Surg* 2017;38:84-9.
167. Tokuda T, Hirano K, Sakamoto Y, Mori S, Kobayashi N, Araki M, et al. Use of the Wound, Ischemia, foot Infection classification system in hemodialysis patients after endovascular treatment for critical limb ischemia. *J Vasc Surg* 2018;67:1762-8.
168. White JV, Rutherford RB, Ryjewski C. Chronic subcritical limb ischemia: a poorly recognized stage of critical limb ischemia. *Semin Vasc Surg* 2007;20:62-7.
169. Salaun P, Desormais I, Lapébie FX, Riviére AB, Aboyans V, Lacroix P, et al. Comparison of ankle pressure, systolic toe pressure, and transcutaneous oxygen pressure to predict major amputation after 1 year in the COPART cohort. *Angiology* 2018. 3319718793566.
170. Graziani L, Silvestro A, Bertone V, Manara E, Andreini R, Sigala A, et al. Vascular involvement in diabetic subjects with ischemic foot ulcer: a new morphologic categorization of disease severity. *Eur J Vasc Endovasc Surg* 2007;33:453-60.
171. Ramanan B, Ahmed A, Wu B, Causey MW, Gasper WJ, Vartanian SM, et al. Determinants of midterm functional outcomes, wound healing, and resources used in a hospital-based limb preservation program. *J Vasc Surg* 2017;66:1765-74.
172. Kobayashi N, Hirano K, Yamawaki M, Araki M, Sakai T, Sakamoto Y, et al. Characteristics and clinical outcomes of repeat endovascular therapy after infrapopliteal balloon angioplasty in patients with critical limb ischemia. *Catheter Cardiovasc Interv* 2018;91:505-14.
173. Leithead C, Novak Z, Spangler E, Passman MA, Witcher A, Patterson MA, et al. Importance of postprocedural Wound, Ischemia, and foot Infection (WIFI) restaging in predicting limb salvage. *J Vasc Surg* 2018;67:498-505.
174. Hirsch AT, Allison MA, Gomes AS, Corriere MA, Duval S, Ershow AG, et al. A call to action: women and peripheral artery disease: a scientific statement from the American Heart Association. *Circulation* 2012;125:1449-72.
175. Marrett E, DiBonaventura M, Zhang Q. Burden of peripheral arterial disease in Europe and the United States: a patient survey. *Health Qual Life Outcomes* 2013;11:175.
176. Howard DP, Banerjee A, Fairhead JF, Hands L, Silver LE, Rothwell PM. Population-based study of incidence, risk factors, outcome, and prognosis of ischemic peripheral arterial events: implications for prevention. *Circulation* 2015;132:1805-15.
177. Olinic DM, Spinu M, Olinic M, Homorodean C, Tataru DA, Liew A, et al. Epidemiology of peripheral artery disease in Europe: VAS Educational Paper. *Int Angiol* 2018;37:327-34.
178. Kojima I, Ninomiya T, Hata J, Fukuhara M, Hirakawa Y, Mukai N, et al. A low ankle brachial index is associated with an increased risk of cardiovascular disease: the Hisayama study. *J Atheroscler Thromb* 2014;21:966-73.
179. Sheng CS, Li Y, Huang QF, Kang YY, Li FK, Wang JG. Pulse waves in the lower extremities as a diagnostic tool of