

76. Rahman A, Ross A, Leighton J *et al*. Double-balloon enteroscopy in Crohn's disease: findings and impact on management in a multicenter retrospective study. *Gastrointest Endosc* 2015;82:102–7.
77. Seiderer J, Herrmann K, Diepolder H *et al*. Double-balloon enteroscopy versus magnetic resonance enteroclysis in diagnosing suspected small-bowel Crohn's disease: results of a pilot study. *Scand J Gastroenterol* 2007;42:1376–85.
78. Kiesslich R, Neurath M.. Surveillance colonoscopy in ulcerative colitis: magnifying chromoendoscopy in the spotlight. *Gut* 2004;53:165–7.
79. Subramanian V, Mannath J, Ragunath K *et al*. Meta-analysis: the diagnostic yield of chromoendoscopy for detecting dysplasia in patients with colonic inflammatory bowel disease. *Aliment Pharmacol Ther* 2011;33:304–12.
80. Iacucci M, Kaplan G, Panaccione R *et al*. A randomized trial comparing high definition colonoscopy alone with high definition dye spraying and electronic virtual chromoendoscopy for detection of colonic neoplastic lesions during IBD surveillance colonoscopy. *Am J Gastroenterol* 2018;113:225–34.
81. N Mohammed, P Kant, Abid F *et al*. High definition white light endoscopy (Hdwle) versus high definition with chromoendoscopy (Hdce) in the detection of dysplasia in long standing ulcerative colitis: a randomized controlled trial (abstract). *Gastrointest Endosc* 2015;81:AB148.
82. Iannone A, Ruospo M, Wong G *et al*. Chromoendoscopy for surveillance in ulcerative colitis and Crohn's disease: a systematic review of randomized trials. *Clin Gastroenterol Hepatol* 2017;15(1684-1697):e11.
83. Iannone A, Ruospo M, Wong G *et al*. Chromoendoscopy for surveillance in ulcerative colitis and Crohn's disease: a systematic review of randomized trials. *Clin Gastroenterol Hepatol* 2016;15:1684–97.
84. Pellise M, Lopez-Ceron M, Rodriguez de Miguel C *et al*. Narrow-band imaging as an alternative to chromoendoscopy for the detection of dysplasia in long-standing inflammatory bowel disease: a prospective, randomized, crossover study. *Gastrointest Endosc* 2011;74:840–8.
85. Ignjatovic A, East JE, Subramanian V *et al*. Narrow band imaging for detection of dysplasia in colitis: a randomized controlled trial. *Am J Gastroenterol* 2012;107:885–90.
86. Marion JF, Wayne JD, Israel Y *et al*. Chromoendoscopy is more effective than standard colonoscopy in detecting dysplasia during long-term surveillance of patients with colitis. *Clin Gastroenterol Hepatol* 2016;14:713–9.
87. Gasia MF, Ghosh S, Panaccione R *et al*. Targeted biopsies identify larger proportions of patients with colonic neoplasia undergoing high-definition colonoscopy, dye chromoendoscopy, or electronic virtual chromoendoscopy. *Clin Gastroenterol Hepatol* 2016;14:704–712.e4.
88. Laine L, Kaltenbach T, Barkun A *et al*. SCENIC international consensus statement on surveillance and management of dysplasia in inflammatory bowel disease. *Gastroenterology* 2015;148:639–51.e28.
89. Mooiweer E, van der Meulen-de Jong AE, Ponsioen CY *et al*. Chromoendoscopy for surveillance in inflammatory bowel disease does not increase neoplasia detection compared with conventional colonoscopy with random biopsies: results from a large retrospective study. *Am J Gastroenterol* 2015;110:1014–21.
90. Moussata D, Allez M, Cazals-Hatem D *et al*. Are random biopsies still useful for the detection of neoplasia in patients with IBD undergoing surveillance colonoscopy with chromoendoscopy? *Gut*; advance online publication, 23 January 2017; pii: gutjnl-2016-311892; doi: 10.1136/gutjnl-2016-311892.
91. Carballal S, Maisterra S, Lopez-Serrano A *et al*. Real-life chromoendoscopy for neoplasia detection and characterisation in long-standing IBD. *Gut* 2018;67:70–8.
92. Samuel S, Bruining D, Loftus E Jr *et al*. Endoscopic skipping of the distal terminal ileum in Crohn's disease can lead to negative results from ileocolonoscopy. *Clin Gastroenterol Hepatol* 2012;10:1253–9.
93. Siddiki H, Fidler J, Fletcher J *et al*. Prospective comparison of state-of-the-art MR enterography and CT enterography in small-bowel Crohn's disease. *AJR Am J Roentgenol* 2009;193:113–21.
94. Buchman A, Miller F, Wallin A *et al*. Videocapsule endoscopy versus barium contrast studies for the diagnosis of Crohn's disease recurrence involving the small intestine. *Am J Gastroenterol* 2004;99:2171–7.
95. Adler J, Punglia D, Dillman J *et al*. Computed tomography enterography findings correlate with tissue inflammation, not fibrosis in resected small bowel Crohn's disease. *Inflamm Bowel Dis* 2012;18:849–56.
96. Church P, Turner D, Feldman B *et al*. Systematic review with meta-analysis: magnetic resonance enterography signs for the detection of inflammation and intestinal damage in Crohn's disease. *Aliment Pharmacol Ther* 2015;41:153–66.
97. Bruining D, Bhatnagar G, Rimola J *et al*. CT and MR enterography in Crohn's disease: current and future applications. *Abdom Imaging* 2015;40:965–74.
98. Deepak P, Fletcher J, Fidler J *et al*. Radiological response is associated with better long-term outcomes and is a potential treatment target in patients with small bowel Crohn's disease. *Am J Gastroenterol* 2016;111:997–1006.
99. Desmond A, O'Regan K, Curran C *et al*. Crohn's disease: factors associated with exposure to high levels of diagnostic radiation. *Gut* 2008;57:1524–9.
100. Chatu S, Subramanian V, Pollok R.. Meta-analysis: diagnostic medical radiation exposure in inflammatory bowel disease. *Aliment Pharmacol Ther* 2012;35:529–39.
101. Silva A, HJ L, Hara A *et al*. Innovations in CT dose reduction strategy: application of the adaptive statistical iterative reconstruction algorithm. *AJR Am J Roentgenol* 2010;194:191–9.
102. Wise P, Schwartz D.. The evaluation and treatment of Crohn perianal fistulae: EUA, EUS, MRI and other imaging modalities. *Gastroenterol Clin N Am* 2012;41:379–91.
103. Schwartz D, Wiersema M, Dudiak K *et al*. A comparison of endoscopic ultrasound, magnetic resonance imaging, and exam under anesthesia for evaluation of Crohn's perianal fistulas. *Gastroenterology* 2001;121:1064–72.
104. Schwartz D, White C, Wise P *et al*. Use of endoscopic ultrasound to guide combination medical and surgical therapy for patients with Crohn's perianal fistulas. *Inflamm Bowel Dis* 2005;11:727–32.
105. Spradlin N, Wise P, Herline A *et al*. A randomized prospective trial of endoscopic ultrasound to guide combination medical and surgical treatment for Crohn's perianal fistulas. *Am J Gastroenterol* 2008;103:2527–35.
106. Villa C, Pompili G, Franceschelli G *et al*. Role of magnetic resonance imaging in evaluation of the activity of perianal Crohn's disease. *Eur J Radiol* 2012;81:616–22.
107. Seastedt K, Trencheva K, Michelassi F *et al*. Accuracy of CT enterography and magnetic resonance enterography imaging to detect lesions preoperatively in patients undergoing surgery for Crohn's disease. *Dis Colon Rectum* 2014;57:1364–70.
108. Xie Y, Zhu W, Li N *et al*. The outcome of initial percutaneous drainage versus surgical drainage for intra-abdominal abscesses in Crohn's disease. *Int J Colorectal Dis* 2012;27:199–206.
109. Melmed G, Siegel C.. Quality improvement in inflammatory bowel disease. *Gastroenterol Hepatol* 2013;9:286–92.
110. Williet N, Sandborn W, Peyrin-Biroulet L.. Patient-reported outcomes as primary end points in clinical trials of inflammatory bowel disease. *Clin Gastro Hep* 2014;12:1246–56.
111. Sauter B, Beglinger C, Girardin M *et al*. Monitoring disease activity and progression in Crohn's disease. A Swiss perspective on the IBD ahead 'optimised monitoring' recommendations. *Digestion* 2014;89:299–309.
112. Van Assche G, Dignass A, Panes J *et al*. The second European evidence-based consensus on the diagnosis and management of Crohn's disease: definitions and diagnosis. *J Crohns Colitis* 2010;4:7–27.
113. Silverberg MS, Satsangi J, Ahmad T *et al*. Toward an integrated clinical, molecular and serological classification of inflammatory bowel disease: report of a Working Party of the 2005 Montreal World Congress of Gastroenterology. *Can J Gastroenterol* 2005;19(Suppl A):5a–36a.
114. Satsangi J, Silverberg M, Vermeire S *et al*. The Montreal classification of inflammatory bowel disease: controversies, consensus, and implications. *Gut* 2006;55:749–53.
115. Levine A, Griffiths A, Markowitz J *et al*. Pediatric modification of the Montreal classification for inflammatory bowel disease: the Paris classification. *Inflamm Bowel Dis* 2011;17:1314–21.
116. Sandborn W, Hanauer S, Van Assche G *et al*. Treating beyond symptoms with a view to improving patient outcomes in inflammatory bowel diseases. *J Crohns Colitis* 2014;8:927–35.
117. Zallot C, Peyrin-Biroulet L.. Deep remission in inflammatory bowel disease: looking beyond symptoms. *Current Gastroenterol Rep* 2013;15:1–7.
118. Peyrin-Biroulet L, WJ S, Sands B *et al*. Selecting therapeutic targets in inflammatory bowel disease (STRIDE): determining therapeutic goals for treat-to-target. *Am J Gastroenterol* 2015;110:1324–8.
119. Klimczak K, Lykowska-Szuber L, Eder P *et al*. The diagnostic usefulness of fecal lactoferrin in the assessment of Crohn's disease activity. *Eur J Intern Med* 2015;26:623–7.
120. Schoepfer A, Beglinger C, Straumann A *et al*. Fecal calprotectin correlates more closely with the Simple Endoscopic Score for Crohn's disease (SES-CD) than CRP, blood leukocytes, and the CDAI. *Am J Gastroenterol* 2010;105:162–9.
121. Boschetti G, Laidet M, Moussata D *et al*. Levels of fecal calprotectin are associated with the severity of postoperative endoscopic recurrence in asymptomatic patients with Crohn's disease. *Am J Gastroenterol* 2015;110:865–72.