

- multispecialty panel using the RAND/UCLA appropriateness method. *Ann Intern Med* 2015; 163:S1–S40.
14. Wang R, Snoey ER, Clements RC, Hern HG, Price D. Effect of head rotation on vascular anatomy of the neck: an ultrasound study. *J Emerg Med* 2006; 31:283–286.
15. O'Grady NP, Alexander M, Burns LA, et al. Guidelines for the prevention of intravascular catheter-related infections. *Clin Infect Dis* 2011; 52:e162–e193.
16. Raad I. Intravascular-catheter-related infections. *Lancet* 1998; 351: 893–898.
17. Gorski L, Hadaway L, Hagle ME, McGoldrick M, Orr M, Doellman D. Infusion therapy standards of practice. *J Infus Nurs* 2016; 39(suppl 1):S65.
18. Moak JH, Lyons MS, Wright SW, Lindsell CJ. Needle and guide-wire visualization in ultrasound-guided internal jugular vein cannulation. *Am J Emerg Med* 2011; 29:432–436.
19. Weekes AJ, Johnson DA, Keller SM, et al. Central vascular catheter placement evaluation using saline flush and bedside echocardiography. *Acad Emerg Med* 2014; 21:65–52.
20. Wilson SP, Assaf S, Lahham S, et al. Simplified point-of-care ultrasound protocol to confirm central venous catheter placement: a prospective study. *World J Emerg Med* 2017; 8:25–28.
21. Weekes AJ, Keller SM, Efune B, Ghali S, Runyon M. Prospective comparison of ultrasound and CXR for confirmation of central vascular catheter placement. *Emerg Med J* 2016; 33:176–180.
22. Cortellaro F, Mellace L, Paglia S, Costantino G, Sher S, Coen D. Contrast enhanced ultrasound vs chest X-ray to determine correct central venous catheter position. *Am J Emerg Med* 2017; 32:78–81.
23. Pittiruti M, La Greca A, Scoppettuolo G. The electrocardiographic method for positioning the tip of central venous catheters. *J Vasc Access* 2011; 12:280–291.
24. Vezzani A, Brusasco C, Palermo S, et al. Ultrasound localization of central vein catheter and detection of postprocedural pneumothorax: an alternative to chest radiography. *Crit Care Med* 2010; 38:533–538.
25. Augoustides JG, Horak J, Ochroch AE, et al. A randomized controlled clinical trial of real-time needle-guided ultrasound for internal jugular venous cannulation in a large university anesthesia department. *J Cardiothorac Vasc Anesth* 2005; 19:310–315.
26. Takatani J, Takeshima M, Okuda K, et al. Equipment enhanced needle visualization: advantages and indications of an ultrasound software package. *Anaesth Intensive Care* 2012; 40:856–860.
27. Sharp R, Cummings M, Fielder A, Mikocka-Walus A, Grech C, Esterman A. The catheter to vein ratio and rates of symptomatic venous thromboembolism in patients with a peripherally inserted central catheter (PICC): a prospective cohort study. *Int J Nurs Stud* 2015; 52:677–685.
28. Hosokawa K, Shime N, Kato Y, Hashimoto S. A randomized trial of ultrasound image-based skin surface marking versus real-time ultrasound-guided internal jugular vein catheterization in infants. *Anesthesiology* 2007; 107:720–724.
29. Karakitsos D, Labropoulos N, De Groot E, et al. Real-time ultrasound-guided catheterisation of the internal jugular vein: a prospective comparison with the landmark technique in critical care patients. *Crit Care* 2006; 10:R162.
30. Brass P, Hellmich M, Kolodziej L, Schick G, Smith AF. Ultrasound guidance versus anatomical landmarks for internal jugular vein catheterization. *Cochrane Database Syst Rev* 2015; 1:CD006962.
31. Rothschild J. Ultrasound guidance of central vein catheterization. In: *Making Health Care Safer*. Rockville, MD; Agency for Healthcare Research and Quality; 2001.
32. Bansal R, Agarwal SK, Tiwari SC, Dash SC. A prospective randomized study to compare ultrasound-guided with non-ultrasound-guided double lumen internal jugular catheter insertion as a temporary hemodialysis access. *Ren Fail* 2005; 27:561–564.
33. Skippen P, Kissoon N. Ultrasound guidance for central vascular access in the pediatric emergency department. *Pediatr Emerg Care* 2007; 23:203–207.
34. Gilbert TB, Seneff MG, Becker RB. Facilitation of internal jugular venous cannulation using an audio-guided Doppler ultrasound vascular access device: results from a prospective, dual-center, randomized, crossover clinical study. *Crit Care Med* 1995; 23:60–65.
35. Milling T, Holden C, Melniker L, et al. Randomized controlled trial of single-operator vs two-operator ultrasound guidance for internal jugular central venous cannulation. *Acad Emerg Med* 2006; 13:245–247.
36. Bellazzini M, Rankin PM, Gangnon RE, Bjoernsen LP. Ultrasound validation of maneuvers to increase internal jugular vein cross-sectional area and decrease compressibility. *Am J Emerg Med* 2009; 27:454–459.
37. Terai C, Anada H, Matsushima S, Shimizu S, Okada Y. Effects of mild Trendelenburg on central hemodynamics and internal jugular vein velocity, cross-sectional area, and flow. *Am J Emerg Med* 1995; 13:255–258.
38. Lewin MR, Stein J, Wang R, et al. Humming is as effective as Valsalva's maneuver and Trendelenburg's position for ultrasonographic visualization of the jugular venous system and common femoral veins. *Ann Emerg Med* 2007; 50:73–77.
39. Blaivas M, Adhikari S. An unseen danger: frequency of posterior vessel wall penetration by needles during attempts to place internal jugular vein central catheters using ultrasound guidance. *Crit Care Med* 2009; 37:2345–2349.
40. Ault MJ, Rosen BT, Ault BW. Inadvertent carotid artery cannulation during ultrasound guided central venous catheterization. *Ann Emerg Med* 2006; 49:721.
41. Mansfield PF, Hohn DC, Fornage BD, Gregurich MA, Ota DM. Complications and failures of subclavian vein catheterizations. *N Engl J Med* 1994; 331:1735–1738.
42. Brass P, Hellmich M, Kolodziej L, Schick G, Smith AF. Ultrasound guidance versus anatomical landmarks for subclavian or femoral vein catheterization. *Cochrane Database Syst Rev* 2015; 1:CD011447.