

CHAPTER 4: MANAGEMENT OF EXACERBATIONS

KEY POINTS:

- An exacerbation of COPD is defined as an event characterized by dyspnea and/or cough and sputum that worsen over < 14 days. Exacerbations of COPD are often associated with increased local and systemic inflammation caused by airway infection, pollution, or other insults to the lungs.
- As the symptoms are not specific to COPD relevant differential diagnoses should be considered, particularly pneumonia, congestive heart failure and pulmonary embolism.
- The goals for treatment of COPD exacerbations are to minimize the negative impact of the current exacerbation and to prevent subsequent events.
- Short-acting inhaled beta₂-agonists, with or without short-acting anticholinergics, are recommended as the initial bronchodilators to treat an exacerbation.
- Maintenance therapy with long-acting bronchodilators should be initiated as soon as possible. In patients with frequent exacerbations and elevated blood eosinophil levels addition of inhaled corticosteroids to the double bronchodilator regimen should be considered.
- In patients with severe exacerbations, systemic corticosteroids can improve lung function (FEV1), oxygenation and shorten recovery time including hospitalization duration. Duration of therapy should not normally be more than 5 days.
- Antibiotics, when indicated, can shorten recovery time, reduce the risk of early relapse, treatment failure, and hospitalization duration. Duration of therapy should be 5 days.
- Methylxanthines are not recommended due to increased side effect profiles.
- Non-invasive mechanical ventilation should be the first mode of ventilation used in COPD patients with acute respiratory failure who have no absolute contraindication because it improves gas exchange, reduces work of breathing and the need for intubation, decreases hospitalization duration and improves survival.
- Exacerbation recovery time varies, taking up to 4-6 weeks to recover, with some patients failing to return to the pre-exacerbation functional state. Following an exacerbation, appropriate measures for exacerbation prevention should be initiated (see **Chapter 3**).

DEFINITION

An exacerbation of chronic obstructive pulmonary disease (ECOPD) is defined as an event characterized by increased dyspnea and/or cough and sputum that worsens in < 14 days which may be accompanied by tachypnea and/or tachycardia and is often associated with increased local and systemic inflammation caused by infection, pollution, or other insult to the airways. ⁽³²⁸⁾

Considerations

Exacerbations of COPD are important events in the management of COPD because they negatively impact health