

77. Longman B, Won N, Aghajanyan A, Sanchez A. PMI IRS country programs: 2021 comparative cost analysis. Rockville, MD. PMI VectorLink Project, Abt Associates Inc. 2022. [Website](#)
78. Gunasekaran K, Sahu SS, Jambulingam P, Das PK. DDT indoor residual spray, still an effective tool to control *Anopheles fluviatilis*-transmitted *Plasmodium falciparum* malaria in India. *Tropical medicine & international health : TM & IH* 2005;10(2):160-8 [Pubmed](#)
79. Stockholm Convention on Persistent Organic Pollutants (POPS). United Nations Environment Programme 2018. [Website](#)
80. Indoor residual spraying : use of indoor residual spraying for scaling up global malaria control and elimination : WHO position statement. Geneva: World Health Organization 2006. [Website](#)
81. WHO Expert Committee on Malaria (1956 ; Athens, Greece). WHO Expert Committee on Malaria, Athens, 20-28 June 1956 : sixth report. World Health Organization. <https://apps.who.int/iris/handle/10665/64537>. Geneva: World Health Organization 1956.
82. Charlwood J.D, Qassim M, Elinsur E.I, Donnelly M, Petrarca V, Billingsley P.F, et al. The impact of indoor residual spraying with malathion on malaria in refugee camps in eastern Sudan. *Acta Tropica* 2001;80(1):1-8 [Pubmed](#) [Journal Website](#)
83. Rowland M, Hewitt S, Durrani N. Prevalence of malaria in Afghan refugee villages in Pakistan sprayed with lambdacyhalothrin or malathion. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1994;88(4):378-9 [Pubmed](#)
84. Wahid S, Stresman GH, Kamal SS, Sepulveda N, Kleinschmidt I, Bousema T, et al. Heterogeneous malaria transmission in long-term Afghan refugee populations: a cross-sectional study in five refugee camps in northern Pakistan. *Malaria journal* 2016;15:245 [Pubmed](#) [Journal](#)
85. Choi L, Pryce J, Garner P. Indoor residual spraying for preventing malaria in communities using insecticide-treated nets. *The Cochrane database of systematic reviews* 2019;(5):CD012688 [Pubmed](#) [Journal Website](#)
86. Pluess B, Tanser FC, Lengeler C, Sharp BL. Indoor residual spraying for preventing malaria. *The Cochrane database of systematic reviews* 2010;(4):CD006657 [Pubmed](#) [Journal](#)
87. WHO Guidance for countries on combining indoor residual spraying and long-lasting insecticidal nets. Geneva: World Health Organization 2014 . [Website](#)
88. Risks associated with scale-back of vector control after malaria transmission has been reduced. Information note. Geneva: World Health Organization 2015. [Website](#)
89. Maia MF, Kliner M, Richardson M, Lengeler C, Moore SJ. Mosquito repellents for malaria prevention. *The Cochrane database of systematic reviews* 2018;(2):CD011595 [Pubmed](#) [Journal Website](#)
90. Gabaldón Figueira JC, Wagah MG, Adipo LB, Wanjiku C, Maia MF. Topical repellents for malaria prevention. *The Cochrane Database of Systematic Reviews* 2023; (8):CD015422 [Journal Website](#)
91. WHO Housing and health guidelines. Geneva: World Health Organization 2018. [Website](#)
92. Keeping the vector out: housing improvements for vector control and sustainable development. Geneva: World Health Organization 2017. [Website](#)
93. Tusting LS, Bottomley C, Gibson H, Kleinschmidt I, Tatem AJ, Lindsay SW, et al. Housing Improvements and Malaria Risk in Sub-Saharan Africa: A Multi-Country Analysis of Survey Data. *PLoS medicine* 2017;14(2):e1002234 [Pubmed](#) [Journal](#)
94. Guyatt HL, Corlett SK, Robinson TP, Ochola SA, Snow RW. Malaria prevention in highland Kenya: indoor residual house-spraying vs. insecticide-treated bednets. *Tropical medicine & international health : TM & IH* 2002;7(4):298-303 [Pubmed](#)
95. Bhatia MR, Fox-Rushby J, Mills A. Cost-effectiveness of malaria control interventions when malaria mortality is low: insecticide-treated nets versus in-house residual spraying in India. *Social science & medicine (1982)* 2004;59(3):525-39 [Pubmed](#)