

33. Grabe M, BM, Bjerklund-Johansen TE, et al. Guidelines on urological infections. EAU Guidelines edition presented at the 25th EAU Annual Congress, Barcelona, 2010 2010.
34. Cai T, Nesi G, Mazzoli S, et al. Asymptomatic bacteriuria treatment is associated with a higher prevalence of antibiotic resistant strains in women with urinary tract infections. *Clin Infect Dis* 2015;61(11):1655–1661.
35. Ronald AR, Nicolle LE, Stamm E, et al. Urinary tract infection in adults: research priorities and strategies. *Int J Antimicrob Agents* 2001;17(4):343–348.
36. Baumgarten DA, Baumgartner BR. Imaging and radiologic management of upper urinary tract infections. *Urol Clin North Am* 1997;24(3):545–569.
37. Pagano MJ, Barbalat Y, Theofanides MC, et al. Diagnostic yield of cystoscopy in the evaluation of recurrent urinary tract infection in women. *Neurourol Urodyn* 2017;36(3):692–696.
38. Yu M, Jia HM, Zhou C, et al. Urinary and fecal metabonomics study of the protective effect of Chaihu-Shu-Gan-San on antibiotic-induced gut microbiota dysbiosis in rats. *Sci Rep* 2017;7:46551.
39. Gupta K, Hooton TM, Naber KG, et al. International clinical practice guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: a 2010 update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *Clin Infect Dis* 2011;52(5):e103–e120.
40. Christiaens TC, De Meyere M, Verschraegen G, et al. Randomised controlled trial of nitrofurantoin versus placebo in the treatment of uncomplicated urinary tract infection in adult women. *Br J Gen Pract* 2002;52(482):729–734.
41. Gágyor I, Bleidorn J, Kochen MM, et al. Ibuprofen versus fosfomycin for uncomplicated urinary tract infection in women: randomised controlled trial. *BMJ* 2015;351:h6544.
42. Bleidorn J, Gágyor I, Kochen MM, et al. Symptomatic treatment (ibuprofen) or antibiotics (ciprofloxacin) for uncomplicated urinary tract infection? Results of a randomized controlled pilot trial. *BMC Med* 2010;8:30.
43. Bleidorn J, Hummers-Pradier E, Schmiemann G, et al. Recurrent urinary tract infections and complications after symptomatic versus antibiotic treatment: follow-up of a randomised controlled trial. *Ger Med Sci* 2016;14:Doc01.
44. Albert X, Huertas I, Pereiro II, et al. Antibiotics for preventing recurrent urinary tract infection in non-pregnant women. *Cochrane Database Syst Rev* 2004;(3):CD001209.
45. Ferry SA, Holm SE, Stenlund H, et al. The natural course of uncomplicated lower urinary tract infection in women illustrated by a randomized placebo controlled study. *Scand J Infect Dis* 2004;36(4):296–301.
46. Vervloet D, Durham S. Adverse reactions to drugs. *BMJ* 1998;316(7143):1511–1514.
47. Bhattacharya S. The facts about penicillin allergy: a review. *J Adv Pharm Technol Res* 2010;1(1):11–17.
48. Boxer MB, Dykewicz MS, Patterson R, et al. The management of patients with sulfonamide allergy. *N Engl J Allergy Proc* 1988;9(3):219–223.
49. Singh N, Gandhi S, McArthur E, et al. Kidney function and the use of nitrofurantoin to treat urinary tract infections in older women. *CMAJ* 2015;187(9):648–656.
50. Nicolle LE. Urinary tract infections in the older adult. *Clin Geriatr Med* 2016;32(3):523–538.
51. Hoang P, Salbu RL. Updated nitrofurantoin recommendations in the elderly: a closer look at the evidence. *Consult Pharm* 2016;31(7):381–384.
52. Kavatha D, Giamarellou H, Alexiou Z, et al. Cefpodoxime-proxetil versus trimethoprim-sulfamethoxazole for short-term therapy of uncomplicated acute cystitis in women. *Antimicrob Agents Chemother* 2003;47(3):897–900.
53. Iravani A, Klimberg I, Briefer C, et al. A trial comparing low-dose, short-course ciprofloxacin and standard 7 day therapy with co-trimoxazole or nitrofurantoin in the treatment of uncomplicated urinary tract infection. *J Antimicrob Chemother* 1999;43(Suppl A):67–75.
54. Arredondo-García JL, Figueroa-Damián R, Rosas A, et al. Comparison of short-term treatment regimen of ciprofloxacin versus long-term treatment regimens of trimethoprim/sulfamethoxazole or norfloxacin for uncomplicated lower urinary tract infections: a randomized, multicentre, open-label, prospective study. *J Antimicrob Chemother* 2004;54(4):840–843.
55. Nicolle LE, Madsen KS, Debeek GO, et al. Three days of pivmecillinam or norfloxacin for treatment of acute uncomplicated urinary infection in women. *Scand J Infect Dis* 2002;34(7):487–492.
56. Henry D, Ellison W, Sullivan J, et al. Treatment of community-acquired acute uncomplicated urinary tract infection with sparflaxacin versus ofloxacin. The Sparflaxacin Multi Center UUTI Study Group. *Antimicrob Agents Chemother* 1998;42(9):2262–2266.
57. Richard GA, Mathew CP, Kirstein JM, et al. Single-dose fluoroquinolone therapy of acute uncomplicated urinary tract infection in women: results from a randomized, double-blind, multicenter trial comparing single-dose to 3-day fluoroquinolone regimens. *Urology* 2002;59(3):334–339.
58. Stein GE. Comparison of single-dose fosfomycin and a 7-day course of nitrofurantoin in female patients with uncomplicated urinary tract infection. *Clin Ther* 1999;21(11):1864–1872.
59. Minassian MA, Lewis DA, Chattopadhyay D, et al. A comparison between single-dose fosfomycin trometamol (Monuril) and a 5-day course of trimethoprim in the treatment of uncomplicated lower urinary tract infection in women. *Int J Antimicrob Agents* 1998;10(1):39–47.
60. Naber KG, Allin DM, Clarysse L, et al. Gatifloxacin 400 mg as a single shot or 200 mg once daily for 3 days is as effective as ciprofloxacin 250 mg twice daily for the treatment of patients with uncomplicated urinary tract infections. *Int J Antimicrob Agents* 2004;23(6):596–605.
61. Fourcroy JL, Berner B, Chiang YK, et al. Efficacy and safety of a novel once-daily extended-release ciprofloxacin tablet formulation for treatment of uncomplicated urinary tract infection in women. *Antimicrob Agents Chemother* 2005;49(10):4137–4143.
62. Henry DC Jr, Bettis RB, Riffer E, et al. Comparison of once-daily extended-release ciprofloxacin and conventional twice-daily ciprofloxacin for the treatment of uncomplicated urinary tract infection in women. *Clin Ther* 2002;24(12):2088–2104.
63. Vogel T, Verreault R, Gourdeau M, et al. Optimal duration of antibiotic therapy for uncomplicated urinary tract infection in older women: a double-blind randomized controlled trial. *CMAJ* 2004;170(4):469–473.
64. Auquer F, Córdón F, Gorina E, et al. Single-dose ciprofloxacin versus 3 days of norfloxacin in uncomplicated urinary tract infections in women. *Clin Microbiol Infect* 2002;8(1):50–54.
65. Hooton TM, Scholes D, Gupta K, et al. Amoxicillin-clavulanate vs ciprofloxacin for the treatment of uncomplicated cystitis in women: a randomized trial. *JAMA* 2005;293(8):949–955.
66. Leigh AP, Nemeth MA, Keyserling CH, et al. Cefdinir versus cefaclor in the treatment of uncomplicated urinary tract infection. *Clin Ther* 2000;22(7):818–825.
67. Huttner A, Verhaegh EM, Harbarth S, et al. Nitrofurantoin revisited: a systematic review and meta-analysis of controlled trials. *J Antimicrob Chemother* 2015;70(9):2456–2464.
68. Garau J. Other antimicrobials of interest in the era of extended-spectrum beta-lactamases: fosfomycin, nitrofurantoin and tigecycline. *Clin Microbiol Infect* 2008;14(Suppl 1):198–202.
69. Seroy JT, Grim SA, Reid GE, et al. Treatment of MDR urinary tract infections with oral fosfomycin: a retrospective analysis. *J Antimicrob Chemother* 2016;71(9):2563–2568.