

35. Corman VM, Landt O, Kaiser M, et al: Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro Surveill* 2020; 25:2000045
36. Chu DKW, Pan Y, Cheng SMS, et al: Molecular diagnosis of a novel coronavirus (2019-nCoV) causing an outbreak of pneumonia. *Clin Chem* 2020; 66:549–555
37. Xie C, Jiang L, Huang G, et al: Comparison of different samples for 2019 novel coronavirus detection by nucleic acid amplification tests. *Int J Infect Dis* 2020; 93:264–267
38. Yam WC, Chan KH, Poon LL, et al: Evaluation of reverse transcription-PCR assays for rapid diagnosis of severe acute respiratory syndrome associated with a novel coronavirus. *J Clin Microbiol* 2003; 41:4521–4524
39. Ai T, Yang Z, Hou H, et al: Correlation of chest CT and RT-PCR testing in coronavirus disease 2019 (COVID-19) in China: A report of 1014 Cases. *Radiology* 2020 Feb 26. [online ahead of print]
40. Wu X, Cai Y, Huang X, et al: Co-infection with SARS-CoV-2 and influenza A virus in patient with pneumonia, China. *Emerg Infect Dis* 2020 Mar 11. [online ahead of print]
41. Chan PK, To WK, Ng KC, et al: Laboratory diagnosis of SARS. *Emerg Infect Dis* 2004; 10:825–831
42. Yang X, Yu Y, Xu J, et al: Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: A single-centered, retrospective, observational study. *Lancet Respir Med* 2020 Feb 24. [online ahead of print]
43. Wang D, Hu B, Hu C, et al: Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020; 323:1061–1069
44. Huang C, Wang Y, Li X, et al: Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; 395:497–506
45. Zhou F, Yu T, Du R, et al: Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: A retrospective cohort study. *Lancet* 2020; 395:1054–1062
46. Ruan Q, Yang K, Wang W, et al: Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med* 2020 Mar 3. [online ahead of print]
47. Bednarczyk JM, Fridfinnson JA, Kumar A, et al: Incorporating dynamic assessment of fluid responsiveness into goal-directed therapy: A systematic review and meta-analysis. *Crit Care Med* 2017; 45:1538–1545
48. Bentzer P, Griesdale DE, Boyd J, et al: Will this hemodynamically unstable patient respond to a bolus of intravenous fluids? *JAMA* 2016; 316:1298–1309
49. Pan J, Peng M, Liao C, et al: Relative efficacy and safety of early lactate clearance-guided therapy resuscitation in patients with sepsis: A meta-analysis. *Medicine (Baltimore)* 2019; 98:e14453
50. Hernandez G, Ospina-Tascon GA, Damiani LP, et al: Effect of a resuscitation strategy targeting peripheral perfusion status vs serum lactate levels on 28-day mortality among patients with septic shock: The ANDROMEDA-SHOCK randomized clinical trial. *JAMA* 2019; 321:654–664
51. Meyhoff TS, Moller MH, Hjortrup PB, et al: Lower versus higher fluid volumes during initial management of sepsis: A systematic review with meta-analysis and trial sequential analysis. *Chest* 2020 Jan 23. [online ahead of print]
52. Silversides JA, Major E, Ferguson AJ, et al: Conservative fluid management or deresuscitation for patients with sepsis or acute respiratory distress syndrome following the resuscitation phase of critical illness: A systematic review and meta-analysis. *Intensive Care Med* 2017; 43:155–170
53. Maitland K, Igili S, Opoka RO, et al: Mortality after fluid bolus in African children with severe infection. *N Engl J Med* 2011; 364:2483–2495
54. Lewis SR, Pritchard MW, Evans DJ, et al: Colloids versus crystalloids for fluid resuscitation in critically ill people. *Cochrane Database Syst Rev* 2018; 8:CD000567
55. Antequera Martin AM, Barea Mendoza JA, Muriel A, et al: Buffered solutions versus 0.9% saline for resuscitation in critically ill adults and children. *Cochrane Database Syst Rev* 2019; 7:CD012247
56. Moller MH, Claudius C, Juntila E, et al: Scandinavian SSAI clinical practice guideline on choice of first-line vasopressor for patients with acute circulatory failure. *Acta Anaesthesiol Scand* 2016; 60:1347–1366
57. Gamper G, Havel C, Arrich J, et al: Vasopressors for hypotensive shock. *Cochrane Database Syst Rev* 2016; 2:CD003709
58. Honarmand K, Um KJ, Belley-Cote EP, et al: Canadian Critical Care Society clinical practice guideline: The use of vasopressin and vasopressin analogues in critically ill adults with distributive shock. *Can J Anaesth* 2020; 67:369–376
59. McIntyre WF, Um KJ, Alhazzani W, et al: Association of vasopressin plus catecholamine vasopressors vs catecholamines alone with atrial fibrillation in patients with distributive shock: A systematic review and meta-analysis. *JAMA* 2018; 319:1889–1900
60. Lamontagne F, Day AG, Meade MO, et al: Pooled analysis of higher versus lower blood pressure targets for vasopressor therapy septic and vasodilatory shock. *Intensive Care Med* 2018; 44:12–21
61. Lamontagne F, Richards-Belle A, Thomas K, et al; 65 Trial Investigators: Effect of reduced exposure to vasopressors on 90-day mortality in older critically ill patients with vasodilatory hypotension: A randomized clinical trial. *JAMA* 2020; 323:938–949
62. Moller MH, Granholm A, Juntila E, et al: Scandinavian SSAI clinical practice guideline on choice of inotropic agent for patients with acute circulatory failure. *Acta Anaesthesiol Scand* 2018; 62:420–450
63. Rygaard SL, Butler E, Granholm A, et al: Low-dose corticosteroids for adult patients with septic shock: A systematic review with meta-analysis and trial sequential analysis. *Intensive Care Med* 2018; 44:1003–1016
64. Lamontagne F, Rochwerg B, Lytvin L, et al: Corticosteroid therapy for sepsis: A clinical practice guideline. *BMJ* 2018; 362:k3284
65. Chen N, Zhou M, Dong X, et al: Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: A descriptive study. *Lancet* 2020; 395:507–513
66. van den Boom W, Hoy M, Sankaran J, et al: The search for optimal oxygen saturation targets in critically ill patients: Observational data from large ICU databases. *Chest* 2020; 157:566–573
67. Chu DK, Kim LH, Young PJ, et al: Mortality and morbidity in acutely ill adults treated with liberal versus conservative oxygen therapy (IOTA): A systematic review and metaanalysis. *Lancet* 2018; 391:1693–1705
68. Siemieniuk RAC, Chu DK, Kim LH, et al: Oxygen therapy for acutely ill medical patients: a clinical practice guideline. *BMJ* 2018; 363:k4169
69. ICU-ROX Investigators and the Australian and New Zealand Intensive Care Society Clinical Trials Group, Mackle D, Bellomo R, et al: Conservative oxygen therapy during mechanical ventilation in the ICU. *N Engl J Med* 2020; 382:989–998
70. Barrot L, Asfar P, Mauny F, et al; LOCO2 Investigators and REVA Research Network: Liberal or conservative oxygen therapy for acute respiratory distress syndrome. *N Engl J Med* 2020; 382:999–1008
71. Frat JP, Thille AW, Mercat A, et al; FLORALI Study Group; REVA Network: High-flow oxygen through nasal cannula in acute hypoxemic respiratory failure. *N Engl J Med* 2015; 372:2185–2196
72. Ni YN, Luo J, Yu H, et al: The effect of high-flow nasal cannula in reducing the mortality and the rate of endotracheal intubation when used before mechanical ventilation compared with conventional oxygen therapy and noninvasive positive pressure ventilation. A systematic review and meta-analysis. *Am J Emerg Med* 2018; 36:226–233
73. Ou X, Hua Y, Liu J, et al: Effect of high-flow nasal cannula oxygen therapy in adults with acute hypoxemic respiratory failure: a meta-analysis of randomized controlled trials. *CMAJ* 2017; 189:E260–E267
74. Rochwerg B, Granton D, Wang DX, et al: High flow nasal cannula compared with conventional oxygen therapy for acute hypoxemic respiratory failure: A systematic review and meta-analysis. *Intensive Care Med* 2019; 45:563–572
75. Raboud J, Shigayeva A, McGeer A, et al: Risk factors for SARS transmission from patients requiring intubation: A multicentre investigation in Toronto, Canada. *PLoS One* 2010; 5:e10717