

- 53 Arnold JB, Mackintosh S, Olds TS, Jones S, Thewlis D. Improvements in knee biomechanics during walking are associated with increased physical activity after total knee arthroplasty. *J Orthop Res*. 2015;33:1818–1825.
- 54 Taniguchi M, Sawano S, Kugo M, Maegawa S, Kawasaki T, Ichihashi N. Physical activity promotes gait improvement in patients with total knee arthroplasty. *J Arthroplasty*. 2016;31:984–988.
- 55 Piva SR, Farrokhi S, Almeida G, Fitzgerald GK, Levison TJ, DiGioia AM. Dose-associated changes in gait parameters in response to exercise programs after total knee arthroplasty: secondary analysis of two randomized studies. *Int J Phys Med Rehabil*. 2015;3:3–7.
- 56 Bruun-Olsen V, Heiberg KE, Wahl AK, Mengshoel AM. The immediate and long-term effects of a walking-skill program compared to usual physiotherapy care in patients who have undergone total knee arthroplasty (TKA): a randomized controlled trial. *Disabil Rehabil*. 2013;35:2008–2015.
- 57 Christiansen CL, Bade MJ, Davidson BS, Dayton MR, Stevens-Lapsley JE. Effects of weight-bearing biofeedback training on functional movement patterns following total knee arthroplasty: a randomized controlled trial. *J Orthop Sports Phys Ther*. 2015;45:647–655.
- 58 Frost H, Lamb SE, Robertson S. A randomized controlled trial of exercise to improve mobility and function after elective knee arthroplasty. Feasibility, results and methodological difficulties. *Clin Rehabil*. 2002;16:200–209.
- 59 Liao CD, Lin LF, Huang YC, Huang SW, Chou LC, Liou TH. Functional outcomes of outpatient balance training following total knee replacement in patients with knee osteoarthritis: a randomized controlled trial. *Clin Rehabil*. 2015;29:855–867.
- 60 Moffet H, Collet JP, Shapiro SH, Paradis G, Marquis F, Roy L. Effectiveness of intensive rehabilitation on functional ability and quality of life after first total knee arthroplasty: a single-blind randomized controlled trial. *Arch Phys Med Rehabil*. 2004;85:546–556.
- 61 Li J, Wu T, Xu Z, Gu X. A pilot study of post-total knee replacement gait rehabilitation using lower limbs robot-assisted training system. *Eur J Orthop Surg Traumatol*. 2014;24:203–208.
- 62 Bin SI, Nam TS. Early results of high-flex total knee arthroplasty: comparison study at 1 year after surgery. *Knee Surg Sports Traumatol Arthrosc*. 2007;15:350–355.
- 63 Cheng K, Ridley D, Bird J, McLeod G. Patients with fixed flexion deformity after total knee arthroplasty do just as well as those without: ten-year prospective data. *Int Orthop*. 2010;34:663–667.
- 64 Gatha NM, Clarke HD, Fuchs R, Scuderi GR, Insall JN. Factors affecting postoperative range of motion after total knee arthroplasty. *J Knee Surg*. 2004;17:196–202.
- 65 Ritter MA, Lutzgring JD, Davis KE, Berend ME, Pierson JL, Meneghini RM. The role of flexion contracture on outcomes in primary total knee arthroplasty. *J Arthroplasty*. 2007;22:1092–1096.
- 66 Ritter MA, Lutzgring JD, Davis KE, Berend ME. The effect of postoperative range of motion on functional activities after posterior cruciate-retaining total knee arthroplasty. *J Bone Joint Surg Am*. 2008;90:777–784.
- 67 Li B, Wang G, Wang Y, Bai L. Effect of two limb positions on venous hemodynamics and hidden blood loss following total knee arthroplasty. *J Knee Surg*. 2017;30:70–74.
- 68 Li B, Wen Y, Liu D, Tian L. The effect of knee position on blood loss and range of motion following total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc*. 2012;20:594–599.
- 69 Panni AS, Cerciello S, Vasso M, Del Regno C. Knee flexion after total knee arthroplasty reduces blood loss. *Knee Surg Sports Traumatol Arthrosc*. 2014;22:1859–1864.
- 70 Yang Y, Yong-Ming L, Pei-jian D, Jia L, Ying-ze Z. Leg position influences early blood loss and functional recovery following total knee arthroplasty: a randomized study. *Int J Surg*. 2015;23(Pt A):82–86.
- 71 De Fine M, Traina F, Giavaresi G, et al. Effect of different postoperative flexion regimes on the outcomes of total knee arthroplasty: randomized controlled trial. *Knee Surg Sports Traumatol Arthrosc*. 2017;25:2972–2977.
- 72 Jiang C, Lou J, Qian W, Ye C, Zhu S. Impact of flexion versus extension of knee position on outcomes after total knee arthroplasty: a meta-analysis. *Arch Orthop Trauma Surg*. 2017;137:257–265.
- 73 Avramidis K, Karachalios T, Popotonasios K, Sacorafas D, Papathanasiades AA, Malizos KN. Does electric stimulation of the vastus medialis muscle influence rehabilitation after total knee replacement? *Orthopedics*. 2011;34:175.
- 74 Avramidis K, Strike PW, Taylor PN, Swain ID. Effectiveness of electric stimulation of the vastus medialis muscle in the rehabilitation of patients after total knee arthroplasty. *Arch Phys Med Rehabil*. 2003;84:1850–1853.
- 75 Stevens-Lapsley JE, Balter JE, Wolfe P, Eckhoff DG, Kohrt WM. Early neuromuscular electrical stimulation to improve quadriceps muscle strength after total knee arthroplasty: a randomized controlled trial. *Phys Ther*. 2012;92:210–226.
- 76 Yoshida Y, Ikuno K, Shomoto K. Comparison of the effect of sensory-level and conventional motor-level neuromuscular electrical stimulations on quadriceps strength after total knee arthroplasty: a prospective randomized single-blind trial. *Arch Phys Med Rehabil*. 2017;98:2364–2370.
- 77 Petterson SC, Mizner RL, Stevens JE, et al. Improved function from progressive strengthening interventions after total knee arthroplasty: a randomized clinical trial with an imbedded prospective cohort. *Arthritis Rheum*. 2009;61:174–183.
- 78 Bade MJ, Struessel T, Dayton M, et al. Early high-intensity versus low-intensity rehabilitation after total knee arthroplasty: a randomized controlled trial. *Arthritis Care Res (Hoboken)*. 2017;69:1360–1368.
- 79 Molla RY, Sadeghi H, Kahlaee AH. The effect of early progressive resistive exercise therapy on balance control of patients with total knee arthroplasty: a randomized controlled trial. *Topics in Geriatric Rehabilitation*. 2017;33:286–294.
- 80 Husby VS, Foss OA, Husby OS, Winther SB. Randomized controlled trial of maximal strength training vs. standard rehabilitation following total knee arthroplasty. *Eur J Phys Rehabil Med*. 2018;54:371–379.
- 81 Lizaur-Utrilla A, Martinez-Mendez D, Miralles-Munoz FA, Marco-Gomez L, Lopez-Prats FA. Comparable outcomes after total knee arthroplasty in patients under 55 years than in older patients: a matched prospective study with minimum follow-up of 10 years. *Knee Surg Sports Traumatol Arthrosc*. 2017;25:3396–3402.
- 82 Naylor JM, Yeo AE, Mittal R, Ko VW, Harris IA. Improvements in knee range and symptomatic and functional behavior after knee arthroplasty based on preoperative restriction in range. *J Arthroplasty*. 2012;27:1100–1105.
- 83 Sullivan M, Tanzer M, Stanish W, et al. Psychological determinants of problematic outcomes following total knee arthroplasty. *Pain*. 2009;143:123–129.
- 84 Xu S, Chen JY, Lo NN, et al. The influence of obesity on functional outcome and quality of life after total knee arthroplasty: a ten-year follow-up study. *Bone Joint J*. 2018;100-b:579–583.
- 85 Liao CD, Huang YC, Lin LF, et al. Continuous passive motion and its effects on knee flexion after total knee arthroplasty in patients with knee osteoarthritis. *Knee Surg Sports Traumatol Arthrosc*. 2016;24:2578–2586.
- 86 Lizaur A, Marco L, Cebrian R. Preoperative factors influencing the range of movement after total knee arthroplasty for severe osteoarthritis. *J Bone Joint Surg Br*. 1997;79:626–629.
- 87 Lizaur-Utrilla A, Miralles-Munoz FA, Sanz-Reig J, Collados-Maestre I. Cementless total knee arthroplasty in