

- obese patients: a prospective matched study with follow-up of 5-10 years. *J Arthroplasty*. 2014;29:1192-1196.
- 88 Rossi MD, Eberle T, Roche M, et al. Use of a squatting movement as a clinical marker of function after total knee arthroplasty. *Am J Phys Med Rehabil*. 2013;92:53-60.
 - 89 Sharma L, Sinacore J, Daugherty C, et al. Prognostic factors for functional outcome of total knee replacement: a prospective study. *J Gerontol A Biol Sci Med Sci*. 1996;51:M152-M157.
 - 90 Si HB, Zeng Y, Zhong J, et al. The effect of primary total knee arthroplasty on the incidence of falls and balance-related functions in patients with osteoarthritis. *Sci Rep*. 2017;7:16583.
 - 91 Stevens-Lapsley JE, Petterson SC, Mizner RL, Snyder-Mackler L. Impact of body mass index on functional performance after total knee arthroplasty. *J Arthroplasty*. 2010;25:1104-1109.
 - 92 Clement ND, Jenkins PJ, MacDonald D, et al. Socioeconomic status affects the Oxford knee score and short-form 12 score following total knee replacement. *Bone Joint J*. 2013;95-b:52-58.
 - 93 Clement ND, MacDonald D, Simpson AH, Burnett R. Total knee replacement in patients with concomitant back pain results in a worse functional outcome and a lower rate of satisfaction. *Bone Joint J*. 2013;95-b:1632-1639.
 - 94 Mizner RL, Petterson SC, Stevens JE, Axe MJ, Snyder-Mackler L. Preoperative quadriceps strength predicts functional ability one year after total knee arthroplasty. *J Rheumatol*. 2005;32:1533-1539.
 - 95 Holm B, Kristensen MT, Bencke J, Husted H, Kehlet H, Bandholm T. Loss of knee-extension strength is related to knee swelling after total knee arthroplasty. *Arch Phys Med Rehabil*. 2010;91:1770-1776.
 - 96 Park KK, Chang CB, Kang YG, Seong SC, Kim TK. Correlation of maximum flexion with clinical outcome after total knee replacement in Asian patients. *J Bone Joint Surg Br*. 2007;89:604-608.
 - 97 Twiggs J, Salmon L, Kolos E, Bogue E, Miles B, Roe J. Measurement of physical activity in the pre- and early post-operative period after total knee arthroplasty for osteoarthritis using a Fitbit flex device. *Med Eng Phys*. 2018;51:31-40.
 - 98 Dauty M, Smitt X, Menu P, Dubois C. Which factors affect the duration of inpatient rehabilitation after total knee arthroplasty in the absence of complications? *Ann Phys Rehabil Med*. 2009;52:234-245.
 - 99 Jones CA, Voaklander DC, Suarez-Alma ME. Determinants of function after total knee arthroplasty. *Phys Ther*. 2003;83:696-706.
 - 100 Kauppila AM, Kyllonen E, Ohtonen P, Leppilahti J, Sintonen H, Arokoski JP. Outcomes of primary total knee arthroplasty: the impact of patient-relevant factors on self-reported function and quality of life. *Disabil Rehabil*. 2011;33:1659-1667.
 - 101 Kramers-de Quervain IA, Kampfen S, Munzinger U, Mannion AF. Prospective study of gait function before and 2 years after total knee arthroplasty. *Knee*. 2012;19:622-627.
 - 102 Tay KS, Cher EWL, Zhang K, Tan SB, Howe TS, Koh JSB. Comorbidities have a greater impact than age alone in the outcomes of octogenarian total knee arthroplasty. *J Arthroplasty*. 2017;32:3373-3378.
 - 103 Collados-Maestre I, Lizaaur-Utrilla A, Martinez-Mendez D, Marco-Gomez L, Lopez-Prats FA. Concomitant low back pain impairs outcomes after primary total knee arthroplasty in patients over 65 years: a prospective, matched cohort study. *Arch Orthop Trauma Surg*. 2016;136:1767-1771.
 - 104 Lin J, Yang B, Weng XS, Jin J, Zhao Q, Qiu GX. Effect of osteoarthritis patients' gender on rehabilitation after total knee arthroplasty. *Chin Med Sci J*. 2009;24:102-106.
 - 105 Hudakova Z, Zieba HR, Lizis P, et al. Evaluation of the effects of a physiotherapy program on quality of life in females after unilateral total knee arthroplasty: a prospective study. *J Phys Ther Sci*. 2016;28:1412-1417.
 - 106 Akbaba YA, Yeldan I, Guney N, Ozdincler AR. Intensive supervision of rehabilitation programme improves balance and functionality in the short term after bilateral total knee arthroplasty. *Knee Surg Sports Traumatol Arthrosc*. 2016;24:26-33.
 - 107 Fransen M, Nairn L, Bridgett L, et al. Post-acute rehabilitation after total knee replacement: a multicenter randomized clinical trial comparing long-term outcomes. *Arthritis Care Res (Hoboken)*. 2017;69:192-200.
 - 108 Artz N, Dixon S, Wylde V, et al. Comparison of group-based outpatient physiotherapy with usual care after total knee replacement: a feasibility study for a randomized controlled trial. *Clin Rehabil*. 2017;31:487-499.
 - 109 Kauppila AM, Kyllonen E, Ohtonen P, et al. Multidisciplinary rehabilitation after primary total knee arthroplasty: a randomized controlled study of its effects on functional capacity and quality of life. *Clin Rehabil*. 2010;24:398-411.
 - 110 Hiyama Y, Kamitani T, Wada O, Mizuno K, Yamada M. Effects of group-based exercise on range of motion, muscle strength, functional ability, and pain during the acute phase after total knee arthroplasty: a controlled clinical trial. *J Orthop Sports Phys Ther*. 2016;46:742-748.
 - 111 Labraca NS, Castro-Sanchez AM, Mataran-Penarrocha GA, Arroyo-Morales M, Sanchez-Joya Mdel M, Moreno-Lorenzo C. Benefits of starting rehabilitation within 24 hours of primary total knee arthroplasty: randomized clinical trial. *Clin Rehabil*. 2011;25:557-566.
 - 112 Pua YH, Ong PH. Association of early ambulation with length of stay and costs in total knee arthroplasty: retrospective cohort study. *Am J Phys Med Rehabil*. 2014;93:962-970.
 - 113 Losina E, Collins JE, Deshpande BR, et al. Financial incentives and health coaching to improve physical activity following total knee replacement: a randomized controlled trial. *Arthritis Care Res (Hoboken)*. 2018;70:732-740.
 - 114 Losina E, Collins JE, Wright J, et al. Postoperative care navigation for total knee arthroplasty patients: a randomized controlled trial. *Arthritis Care Res (Hoboken)*. 2016;68:1252-1259.
 - 115 McLawhorn AS, Fu MC, Schairer WW, Sculco PK, MacLean CH, Padgett DE. Continued inpatient care after primary total knee arthroplasty increases 30-day post-discharge complications: a propensity score-adjusted analysis. *J Arthroplasty*. 2017;32:S113-S118.
 - 116 Padgett DE, Christ AB, Joseph AD, Lee YY, Haas SB, Lyman S. Discharge to inpatient rehab does not result in improved functional outcomes following primary total knee arthroplasty. *J Arthroplasty*. 2018;33:1663-1667.
 - 117 Tribe KL, Lapsley HM, Cross MJ, Courtenay BG, Brooks PM, March LM. Selection of patients for inpatient rehabilitation or direct home discharge following total joint replacement surgery: a comparison of health status and out-of-pocket expenditure of patients undergoing hip and knee arthroplasty for osteoarthritis. *Chronic Illn*. 2005;1:289-302.
 - 118 Brennan GP, Fritz JM, Houck LT, Hunter SJ. Outpatient rehabilitation care process factors and clinical outcomes among patients discharged home following unilateral total knee arthroplasty. *J Arthroplasty*. 2015;30:885-890.
 - 119 Smith BA, Fields CJ, Fernandez N. Physical therapists make accurate and appropriate discharge recommendations for patients who are acutely ill. *Phys Ther*. 2010;90:693-703.
 - 120 Falvey JR, Burke RE, Malone D, Ridgeway KJ, McManus BM, Stevens-Lapsley JE. Role of physical therapists in reducing hospital readmissions: optimizing outcomes for older adults during care transitions from hospital to community. *Phys Ther*. 2016;96:1125-1134.