

- multi-centre retrospective propensity-weighted study. *Int J Antimicrob Agents* **2023**; 62:106809.
198. Nukaga M, Papp-Wallace KM, Hoshino T, et al. Probing the mechanism of inactivation of the FOX-4 cephalomycinase by avibactam. *Antimicrob Agents Chemother* **2018**; 62:e02371-17.
199. Cheng L, Nelson BC, Mehta M, et al. Piperacillin-tazobactam versus other antibacterial agents for treatment of bloodstream infections due to AmpC beta-lactamase-producing enterobacteriaceae. *Antimicrob Agents Chemother* **2017**; 61:e00276-17.
200. Lu B, Wong M, Ha D, et al. Piperacillin/tazobactam versus cefepime or carbapenems for cefoxitin-non-susceptible *Enterobacter cloacae*, *Klebsiella aerogenes*, *Citrobacter freundii*, *Serratia marcescens* and *Morganella morganii* bacteraemia in immunocompromised patients. *J Antimicrob Chemother* **2023**; 78:1009–14.
201. Stewart AG, Paterson DL, Young B, et al. Meropenem versus piperacillin-tazobactam for definitive treatment of bloodstream infections caused by AmpC beta-lactamase-producing *Enterobacter* spp, *Citrobacter freundii*, *Morganella morganii*, *Providencia* spp, or *Serratia marcescens*: a pilot multicenter randomized controlled trial (MERINO-2). *Open Forum Infect Dis* **2021**; 8: ofab387.
202. Zhanel GG, Lawrence CK, Adam H, et al. Imipenem-relebactam and meropenem-vaborbactam: two novel carbapenem-beta-lactamase inhibitor combinations. *Drugs* **2018**; 78:65–98.
203. Tselepis L, Langley GW, Aboklaish AF, et al. In vitro efficacy of imipenem-relebactam and cefepime-AAI101 against a global collection of ESBL-positive and carbapenemase-producing Enterobacteriaceae. *Int J Antimicrob Agents* **2020**; 56:105925.
204. Sader HS, Mendes RE, Doyle TB, Davis AP, Castanheira M. Characterization of *Enterobacter cloacae* and *Citrobacter freundii* species complex isolates with decreased susceptibility to cephalosporins from United States hospitals and activity of ceftazidime/avibactam and comparator agents. *JAC Antimicrob Resist* **2021**; 3:dlab136.
205. Torres A, Wible M, Tawadrous M, et al. Efficacy and safety of ceftazidime/avibactam in patients with infections caused by beta-lactamase-producing Gram-negative pathogens: a pooled analysis from the phase 3 clinical trial programme. *J Antimicrob Chemother* **2023**; 78:2672–82.
206. Shields RK, Iovleva A, Kline EG, Kawai A, McElheny CL, Doi Y. Clinical evolution of AmpC-mediated ceftazidime-avibactam and cefiderocol resistance in *Enterobacter cloacae* complex following exposure to cefepime. *Clin Infect Dis* **2020**; 71:2713–16.
207. Kawai A, McElheny CL, Iovleva A, et al. Structural basis of reduced susceptibility to ceftazidime-avibactam and cefiderocol in *Enterobacter cloacae* due to AmpC R2 loop deletion. *Antimicrob Agents Chemother* **2020**; 64:e00198-20.
208. Shropshire WC, Endres BT, Borjan J, et al. High-level ceftazidime/avibactam resistance in *Escherichia coli* conferred by the novel plasmid-mediated beta-lactamase CMY-185 variant. *J Antimicrob Chemother* **2023**; 78:2442–50.
209. Golden AR, Adam HJ, Baxter M, et al. In vitro activity of cefiderocol, a novel siderophore cephalosporin, against gram-negative bacilli isolated from patients in Canadian intensive care units. *Diagn Microbiol Infect Dis* **2020**; 97:115012.
210. Tato M, Garcia-Castillo M, Bofarull AM, Canton R, Group CS. In vitro activity of ceftolozane/tazobactam against clinical isolates of *Pseudomonas aeruginosa* and Enterobacteriaceae recovered in Spanish medical centres: results of the CENIT study. *Int J Antimicrob Agents* **2015**; 46:502–10.
211. Robin F, Auzou M, Bonnet R, et al. In vitro activity of ceftolozane-tazobactam against *Enterobacter cloacae* complex clinical isolates with different beta-lactam resistance phenotypes. *Antimicrob Agents Chemother* **2018**; 62:e00675-18.
212. Stewart AG, Harris PNA, Chatfield MD, Littleford R, Paterson DL. Ceftolozane-tazobactam versus meropenem for definitive treatment of bloodstream infection due to extended-spectrum beta-lactamase (ESBL) and AmpC-producing Enterobacteriales (“MERINO-3”): study protocol for a multicentre, open-label randomised non-inferiority trial. *Trials* **2021**; 22:301.
213. Fox MT, Melia MT, Same RG, Conley AT, Tamma PD. A seven-day course of TMP-SMX may be as effective as a seven-day course of ciprofloxacin for the treatment of pyelonephritis. *Am J Med* **2017**; 130:842–45.
214. Centers for Disease Control and Prevention. About carbapenem resistant Enterobacteriales. Available at: https://www.cdc.gov/cre/about/?CDC_AAref_Val=https://www.cdc.gov/hai/organisms/cre/index.html. Accessed 24 August 2024.
215. Shropshire WC, Aitken SL, Pifer R, et al. IS26-mediated amplification of blaOXA-1 and blaCTX-M-15 with concurrent outer membrane porin disruption associated with de novo carbapenem resistance in a recurrent bacteraemia cohort. *J Antimicrob Chemother* **2021**; 76:385–95.
216. Sabour S, Huang Y, Bhatnagar A, et al. Detection and characterization of targeted carbapenem-resistant healthcare-associated threats: findings from the antibiotic resistance laboratory network, 2017 to 2019. *Antimicrob Agents Chemother* **2021**; 65:e0110521.
217. van Duin D, Arias CA, Komarow L, et al. Molecular and clinical epidemiology of carbapenem-resistant Enterobacteriales in the USA (CRACKLE-2): a prospective cohort study. *Lancet Infect Dis* **2020**; 20:731–41.
218. Sader HS, Mendes RE, Carvalhaes CG, Kimbrough JH, Castanheira M. Changing epidemiology of carbapenemases among carbapenem-resistant Enterobacteriales from United States hospitals and the activity of aztreonam-avibactam against contemporary enterobacteriales (2019–2021). *Open Forum Infect Dis* **2023**; 10: ofad046.
219. Aitken SL, Tarrand JJ, Deshpande LM, et al. High rates of nonsusceptibility to ceftazidime-avibactam and identification of New Delhi metallo-beta-lactamase production in enterobacteriaceae bloodstream infections at a major cancer center. *Clin Infect Dis* **2016**; 63:954–58.
220. Senchyna F, Gaur RL, Sandlund J, et al. Diversity of resistance mechanisms in carbapenem-resistant Enterobacteriales at a health care system in Northern California, from 2013 to 2016. *Diagn Microbiol Infect Dis* **2019**; 93:250–7.
221. Mojica MF, Rossi MA, Vila AJ, Bonomo RA. The urgent need for metallo-beta-lactamase inhibitors: an unattended global threat. *Lancet Infect Dis* **2022**; 22:e28–34.
222. Tamma PD, Simner PJ. Phenotypic detection of carbapenemase-producing organisms from clinical isolates. *J Clin Microbiol* **2018**; 56:e01140-18.
223. Sorli L, Luque S, Li J, et al. Colistin for the treatment of urinary tract infections caused by extremely drug-resistant *Pseudomonas aeruginosa*: dose is critical. *J Infect* **2019**; 79:253–61.
224. Sandri AM, Landersdorfer CB, Jacob J, et al. Population pharmacokinetics of intravenous polymyxin B in critically ill patients: implications for selection of dosage regimens. *Clin Infect Dis* **2013**; 57:524–31.
225. Satlin MJ, Kubin CJ, Blumenthal JS, et al. Comparative effectiveness of aminoglycosides, polymyxin B, and tigecycline for clearance of carbapenem-resistant *Klebsiella pneumoniae* from urine. *Antimicrob Agents Chemother* **2011**; 55: 5893–9.
226. Wagenlehner FM, Sobel JD, Newell P, et al. Ceftazidime-avibactam versus doripenem for the treatment of complicated urinary tract infections, including acute pyelonephritis: RECAPTURE, a phase 3 randomized trial program. *Clin Infect Dis* **2016**; 63:754–62.
227. Kaye KS, Bhowmick T, Metallidis S, et al. Effect of meropenem-vaborbactam vs piperacillin-tazobactam on clinical cure or improvement and microbial eradication in complicated urinary tract infection: the TANGO I randomized clinical trial. *JAMA* **2018**; 319:788–99.
228. Portsmouth S, van Veenhuyzen D, Echols R, et al. Cefiderocol versus imipenem-cilastatin for the treatment of complicated urinary tract infections caused by Gram-negative uropathogens: a phase 2, randomised, double-blind, non-inferiority trial. *Lancet Infect Dis* **2018**; 18:1319–28.
229. Sims M, Mariyanovski V, McLeroth P, et al. Prospective, randomized, double-blind, phase 2 dose-ranging study comparing efficacy and safety of imipenem/cilastatin plus relebactam with imipenem/cilastatin alone in patients with complicated urinary tract infections. *J Antimicrob Chemother* **2017**; 72:2616–26.
230. Bassetti M, Echols R, Matsunaga Y, et al. Efficacy and safety of cefiderocol or best available therapy for the treatment of serious infections caused by carbapenem-resistant Gram-negative bacteria (CREDIBLE-CR): a randomised, open-label, multicentre, pathogen-focused, descriptive, phase 3 trial. *Lancet Infect Dis* **2021**; 21:226–40.
231. USCAST, The National Antimicrobial Susceptibility Testing Committee for the United States. Aminoglycoside In Vitro Susceptibility Test Interpretive Criteria Evaluations. Version 1.3, 2019. Available at: <http://www.uscast.org>
232. Adelman MW, Bower CW, Grass JE, et al. Distinctive features of ertapenem-mono-resistant carbapenem-resistant Enterobacteriales in the United States: a cohort study. *Open Forum Infect Dis* **2022**; 9:ofab643.
233. Daikos GL, Tsakouli S, Tzouveleki LS, et al. Carbapenemase-producing *Klebsiella pneumoniae* bloodstream infections: lowering mortality by antibiotic combination schemes and the role of carbapenems. *Antimicrob Agents Chemother* **2014**; 58:2322–8.
234. Roberts JA, Kirkpatrick CM, Roberts MS, Robertson TA, Dalley AJ, Lipman J. Meropenem dosing in critically ill patients with sepsis and without renal dysfunction: intermittent bolus versus continuous administration? Monte Carlo dosing simulations and subcutaneous tissue distribution. *J Antimicrob Chemother* **2009**; 64:142–50.
235. Kuti JL, Dandekar PK, Nightingale CH, Nicolau DP. Use of Monte Carlo simulation to design an optimized pharmacodynamic dosing strategy for meropenem. *J Clin Pharmacol* **2003**; 43:1116–23.
236. Li C, Kuti JL, Nightingale CH, Nicolau DP. Population pharmacokinetic analysis and dosing regimen optimization of meropenem in adult patients. *J Clin Pharmacol* **2006**; 46:1171–8.