

- after surgery (the VIGILANCE pilot study)—a randomized controlled pilot trial. *Pilot Feasibility Stud* 2019; 5:36
14. Downey C, Randell R, Brown J, et al: Continuous versus intermittent vital signs monitoring in patients admitted to surgical wards: A cluster-randomised, controlled trial. *Colorectal Dis* 2018; 20(Supplement 4):60
 15. Xie L, Garg T, Svec D, et al: Reducing telemetry use is safe: A retrospective analysis of rapid response team and code events after a successful intervention to reduce telemetry use. *Am J Med Qual* 2019; 34:398–401
 16. Barwise A, Thongprayoon C, Herasevich V, et al: The association between monitoring and RRT activation. *Crit Care Med* 2015; 43(12 SUPPL. 1):51
 17. Subbe CP, Duller B, Bellomo R: Effect of an automated notification system for deteriorating ward patients on clinical outcomes. *Crit Care* 2017; 21:52
 18. Scott Evans R, Kuttler KG, Simpson KJ, et al: Automated detection of physiologic deterioration in hospitalized patients. *J Am Med Inform Assoc* 2015; 22:350–360
 19. Heller AR, Mees ST, Lauterwald B, et al: Detection of deteriorating patients on surgical wards outside the ICU by an automated MEWS-based early warning system with paging functionality. *Ann Surg* 2020; 271:100–105
 20. Bellomo R, Ackerman M, Bailey M, et al: Vital Signs to Identify, Target, and Assess Level of Care Study (VITAL Care Study) Investigators: A controlled trial of electronic automated advisory vital signs monitoring in general hospital wards. *Crit Care Med* 2012; 40:2349–2361
 21. Zimlichman E, Terrence J, Argaman D, et al: Effect of contactless continuous patient monitoring in a medical-surgical unit on intensive care unit transfers: A controlled clinical trial. *Am J Respir Crit Care Med* 2012; 185:A3793
 22. Long C, Murray K, Sparkes C: The use of electronic patient observations tracking and targeted critical care outreach to reduce in-hospital cardiac arrests. *J Intensive Care Soc* 2016; 17(4 Supplement 1):133
 23. Weller RS, Foard KL, Harwood TN: Evaluation of a wireless, portable, wearable multi-parameter vital signs monitor in hospitalized neurological and neurosurgical patients. *J Clin Monit Comput* 2018; 32:945–951
 24. Brown H, Terrence J, Vasquez P, et al: Continuous monitoring in an inpatient medical-surgical unit: A controlled clinical trial. *Am J Med* 2014; 127:226–232
 25. Verrillo SC, Cvach M, Hudson KW, et al: Using continuous vital sign monitoring to detect early deterioration in adult postoperative inpatients. *J Nurs Care Qual* 2019; 34:107–113
 26. Wiles MD, El-Nayal A, Elton G, et al: The effect of patient ethnicity on the accuracy of peripheral pulse oximetry in patients with COVID-19 pneumonitis: A single-centre, retrospective analysis. *Anaesthesia* 2022; 77:143–152
 27. Knight MJ, Subbe CP, Inada-Kim M: Racial discrepancies in oximetry: Where do we stand? *Anaesthesia* 2022; 77:129–131
 28. Hillman K, Chen J, Cretikos M, et al: MERIT study investigators: Introduction of the medical emergency team (MET) system: A cluster-randomised controlled trial. *Lancet* 2005; 365:2091–2097
 29. Davis DP, Aguilar SA, Graham PG, et al: A novel configuration of a traditional rapid response team decreases non-intensive care unit arrests and overall hospital mortality. *J Hosp Med* 2015; 10:352–357
 30. Chatwood AK, Osborne M: The impact of an integrated resuscitation and critical care training programme on cardiac arrest rates and survival after cardiac arrest. *Resuscitation* 2010; 81:S96
 31. Buist MD, Moore GE, Bernard SA, et al: Effects of a medical emergency team on reduction of incidence of and mortality from unexpected cardiac arrests in hospital: Preliminary study. *BMJ* 2002; 324:387–390
 32. Bradley P, DePolo M, Tipton J, et al: Improving sepsis survival using mews for early recognition and immediate response to patient decline. *Crit Care Med* 2019; 47(1 Supplement 1):788
 33. Bunkenborg G, Samuelson K, Poulsen I, et al: Lower incidence of unexpected in-hospital death after interprofessional implementation of a bedside track-and-trigger system. *Resuscitation* 2014; 85:424–430
 34. Campello G, Granja C, Carvalho F, et al: Immediate and long-term impact of medical emergency teams on cardiac arrest prevalence and mortality: A plea for periodic basic life-support training programs. *Crit Care Med* 2009; 37:3054–3061
 35. Fuhrmann L, Perner A, Klausen TW, et al: The effect of multi-professional education on the recognition and outcome of patients at risk on general wards. *Resuscitation* 2009; 80:1357–1360
 36. Guirgis FW, Jones L, Esma R, et al: Managing sepsis: Electronic recognition, rapid response teams, and standardized care save lives. *J Crit Care* 2017; 40:296–302
 37. Heal M, Silvest-Guerrero S, Kohtz C: Design and development of a proactive rapid response team. *Computers Inform Nurs* 2017; 35:77–83
 38. Jones D, Bates S, Warrillow S, et al: Effect of an education programme on the utilization of a medical emergency team in a teaching hospital. *Intern Med J* 2006; 36:231–236
 39. Jones D, Opdam H, Egi M, et al: Long-term effect of a medical emergency team on mortality in a teaching hospital. *Resuscitation* 2007; 74:235–241
 40. Joshi KS, Campbell VK, Gooch RJ, et al: Adult deterioration detection system (Q-ADDS) based rapid response system (RRS) reduces severity of illness and length of stay of ICU admissions from the ward in a regional hospital. *Intensive Care Med Exp* 2015; 3(Supplement 1):A141
 41. Leech C, Cosgrove J, Laws P: MEWS (Modified Early Warning Score) chart completion as a marker for quality in the early recognition of acutely ill patients. The experience of a UK University Teaching Hospital NHS Trust. *J Intensive Care Soc* 2014; 15(1 SUPPL. 1):S32
 42. Lighthall GK, Parast LM, Rapoport L, et al: Introduction of a rapid response system at a United States Veterans Affairs hospital reduced cardiac arrests. *Anesth Analg* 2010; 111:679–686
 43. Merriel A, van der Nelson H, Merriel S, et al: Identifying deteriorating patients through multidisciplinary team training. *Am J Med Qual* 2016; 31:589–595
 44. Mullany DV, Ziegenfuss M, Goleby MA, et al: Improved hospital mortality with a low MET dose: The importance of a modified early warning score and communication tool. *Anaesth Intensive Care* 2016; 44:734–741