

18. Fogas J, Kiss K, Gyura F, et al: Effects of proton pump inhibitor versus H2-receptor antagonist stress ulcer prophylaxis on ventilator-associated pneumonia: A pilot study. *Crit Care* 2013; 17:P402
19. Hata M, Shiono M, Sekino H, et al: Prospective randomized trial for optimal prophylactic treatment of the upper gastrointestinal complications after open heart surgery. *Circ J* 2005; 69:331–334
20. Kantorova I, Svoboda P, Scheer P, et al: Stress ulcer prophylaxis in critically ill patients: A randomized controlled trial. *Hepatogastroenterology* 2004; 51:757–761
21. Kotlyanskaya A, Luka B, Mukherji R: A comparison of lansoprazole disintegrating tablet, lansoprazole suspension or ranitidine for stress ulcer prophylaxis in critically ill patients. *Crit Care Med* 2008; 7:A194
22. Lee T-H, Hung F-M, Yang L-H: Comparison of the efficacy of esomeprazole and famotidine against stress ulcers in a neurosurgical intensive care unit. *Adv Diges Med* 2014; 1:50–53
23. Levy MJ, Seelig CB, Robinson NJ, et al: Comparison of omeprazole and ranitidine for stress ulcer prophylaxis. *Dig Dis Sci* 1997; 42:1255–1259
24. Lou W, Xia Y, Xiang P, et al: Prevention of upper gastrointestinal bleeding in critically ill Chinese patients: A randomized, double-blind study evaluating esomeprazole and cimetidine. *Curr Med Res Opin* 2018; 34:1449–1455
25. Risaliti A, Terrosu A, Uzzau R, et al: Intravenous omeprazole vs ranitidine in the prophylaxis of stress ulcers. *Acta Chir Ital* 1993; 49:397–401
26. Solouki M, Marashian SM, Kouchak M, et al: Comparison between the preventive effects of ranitidine and omeprazole on upper gastrointestinal bleeding among ICU patients. *Tanaffos* 2009; 8:37–42
27. Somberg L, Morris J, Fantus R, et al: Intermittent intravenous pantoprazole and continuous cimetidine infusion: Effect on gastric pH control in critically ill patients at risk of developing stress-related mucosal disease. *J Trauma* 2008; 64:1202–1210
28. Wee B, Liu CH, Cohen H, et al: 731: IV famotidine vs. IV pantoprazole for stress ulcer prevention in the ICU: A prospective study. *Crit Care Med* 2013; 41:A181
29. Terzi CC, Dragosavac D, Coelho NJ, et al: Ranitidine is unable to maintain gastric pH levels above 4 in septic patients. *J Crit Care* 2009; 24:627.e7–627.13
30. Selvanderan S, Summers M, Finnis M, et al: Pantoprazole or Placebo for Stress Ulcer Prophylaxis (POP-UP): Randomized double-blind exploratory study. *Crit Care Med* 2016; 44:1842–1850
31. Alhazzani W, Guyatt G, Alshahrani M, et al; Canadian Critical Care Trials Group: Withholding pantoprazole for stress ulcer prophylaxis in critically ill patients: A pilot randomized clinical trial and meta-analysis. *Crit Care Med* 2017; 45:1121–1129
32. Krag M, Marker S, Perner A, et al; SUP-ICU trial group: Pantoprazole in patients at risk for gastrointestinal bleeding in the ICU. *N Engl J Med* 2018; 379:2199–2208
33. Lin C-C, Hsu Y-L, Chung C-S, et al: Stress ulcer prophylaxis in patients being weaned from the ventilator in a respiratory care center: A randomized control trial. *J Form Med Assoc* 2016; 115:19–24
34. El-Kersh K, Jalil B, McClave SA, et al: Enteral nutrition as stress ulcer prophylaxis in critically ill patients: A randomized controlled exploratory study. *J Crit Care* 2018; 43:108–113
35. Wang Y, Ge L, Ye Z, et al: Efficacy and safety of gastrointestinal bleeding prophylaxis in critically ill patients: An updated systematic review and network meta-analysis of randomized trials. *Intensive Care Med* 2020; 46:1987–2000
36. Reynolds PM, MacLaren R: Re-evaluating the utility of stress ulcer prophylaxis in the critically ill patient: A clinical scenario-based meta-analysis. *Pharmacotherapy* 2019; 39:408–420
37. Liu B, Liu S, Yin A, et al: Risks and benefits of stress ulcer prophylaxis in adult neurocritical care patients: A systematic review and meta-analysis of randomized controlled trials. *Crit Care* 2015; 19:409
38. Daou M, Dionne JC, Teng JF, et al: Prophylactic acid suppressants in patients with primary neurologic injury: A systematic review and meta-analysis of randomized studies. *J Crit Care* 2022; 71:154093
39. Huang HB, Jiang W, Wang CY, et al: Stress ulcer prophylaxis in intensive care unit patients receiving enteral nutrition: A systematic review and meta-analysis. *Crit Care* 2018; 22:20
40. He N, Yan Y, Su S, et al: Are proton pump inhibitors more effective than histamine-2-receptor antagonists for stress ulcer prophylaxis in critically ill patients? A systematic review and meta-analysis of cohort studies. *Ann Pharmacother* 2022; 56:988–997
41. Lee TC, Goodwin Wilson M, Lawandi A, et al: Proton pump inhibitors versus histamine-2 receptor antagonists likely increase mortality in critical care: An updated meta-analysis. *Am J Med* 2021; 134:e184–e188
42. Deliwala SS, Hamid K, Goyal H, et al: Proton pump inhibitors versus histamine-2-receptor antagonists for stress ulcer prophylaxis in critically ill patients: A meta-analysis and trial sequential analysis. *J Clin Gastroenterol* 2022; 56:204–217
43. Reynolds PM, Wells L, Powell M, et al: Associated mortality risk of proton pump inhibitor therapy for the prevention of stress ulceration in intensive care unit patients: A systematic review and meta-analysis. *J Clin Gastroenterol* 2022; 57:586–594
44. Farrell CP, Mercogliano G, Kuntz CL: Overuse of stress ulcer prophylaxis in the critical care setting and beyond. *J Crit Care* 2010; 25:214–220
45. Domond M, Struby C, Berson A, et al: The downward spiral of acid suppressive therapy in stress ulcer prophylaxis. *Am J Gastroenterol* 2017; 112:S621–S622
46. Boyd C, Hassig T, MacLaren R: A pragmatic assessment of proton pump inhibitors vs. histamine type 2 receptor antagonists on clinically important gastrointestinal bleeding and mortality when used for stress ulcer prophylaxis in the ICU. *Pharmacotherapy* 2021; 41:820–827
47. MacLaren R, Reynolds PM, Allen RR: Histamine-2 receptor antagonists vs proton pump inhibitors on gastrointestinal tract hemorrhage and infectious complications in the intensive care unit. *JAMA Intern Med* 2014; 174:564–574
48. Marker S, Perner A, Wetterslev J, et al; SUP-ICU investigators: Pantoprazole prophylaxis in ICU patients with high severity of disease: A post hoc analysis of the placebo-controlled SUP-ICU trial. *Intensive Care Med* 2019; 45:609–618
49. Murphy CE, Stevens AM, Ferrentino N, et al: Frequency of inappropriate continuation of acid suppressive therapy after discharge in patients who began therapy in the surgical intensive care unit. *Pharmacotherapy* 2008; 28:968–976