

- [83] Inohara T, Kohsaka S, Fukuda K, Menon V. The challenges in the management of right ventricular infarction. *Eur Heart J Acute Cardiovasc Care*. 2013;2:226–34.
- [84] Haji SA, Movahed A. Right ventricular infarction–diagnosis and treatment. *Clin Cardiol*. 2000;23:473–82.
- [85] de Winter RW, Adams R, Verouden NJ, de Winter RJ. Precordial junctional ST-segment depression with tall symmetric T-waves signifying proximal LAD occlusion, case reports of STEMI equivalence. *J Electrocardiol*. 2016;49:76–80.
- [86] Smith SW, Dodd KW, Henry TD, Dvorak DM, Pearce LA. Diagnosis of ST-elevation myocardial infarction in the presence of left bundle branch block with the ST-elevation to S-wave ratio in a modified Sgarbossa rule. *Ann Emerg Med*. 2012;60:766–76.
- [87] Cai Q, Mehta N, Sgarbossa EB, Pinski SL, Wagner GS, Califf RM, et al. The left bundle-branch block puzzle in the 2013 ST-elevation myocardial infarction guideline: from falsely declaring emergency to denying reperfusion in a high-risk population. Are the Sgarbossa Criteria ready for prime time? *Am Heart J*. 2013;166:409–13.
- [88] Sgarbossa EB, Pinski SL, Barbagelata A, Underwood DA, Gates KB, Topol EJ, et al. Electrocardiographic diagnosis of evolving acute myocardial infarction in the presence of left bundle-branch block. GUSTO-1 (Global Utilization of Streptokinase and Tissue Plasminogen Activator for Occluded Coronary Arteries) Investigators. *N Engl J Med*. 1996;334:481–7.
- [89] de Zwaan C, Bär FW, Wellens HJ. Characteristic electrocardiographic pattern indicating a critical stenosis high in left anterior descending coronary artery in patients admitted because of impending myocardial infarction. *Am Heart J*. 1982;103:730–6.
- [90] Knotts RJ, Wilson JM, Kim E, Huang HD, Birnbaum Y. Diffuse ST depression with ST elevation in aVR: Is this pattern specific for global ischemia due to left main coronary artery disease? *J Electrocardiol*. 2013;46:240–8.
- [91] Wong CK, Gao W, Stewart RA, Benatar J, French JK, Aylward PE, et al. aVR ST elevation: an important but neglected sign in ST elevation acute myocardial infarction. *Eur Heart J*. 2010;31:1845–53.
- [92] Koechlin L, Strebel I, Zimmermann T, Nestelberger T, Walter J, Lopez-Ayala P, et al. Hyperacute T wave in the early diagnosis of acute myocardial infarction. *Ann Emerg Med*. 2023;82:194–202.
- [93] Smith SW, Meyers HP. Hyperacute T-waves can be a useful sign of occlusion myocardial infarction if appropriately defined. *Ann Emerg Med*. 2023;82:203–6.
- [94] Kaul P, Fu Y, Chang WC, Harrington RA, Wagner GS, Goodman SG, et al. Prognostic value of ST segment depression in acute coronary syndromes: insights from PARAGON-A applied to GUSTO-IIb. PARAGON-A and GUSTO IIb Investigators. Platelet IIb/IIIa Antagonism for the Reduction of Acute Global Organization Network. *J Am Coll Cardiol*. 2001;38:64–71.
- [95] Savonitto S, Cohen MG, Politi A, Hudson MP, Kong DF, Huang Y, et al. Extent of ST-segment depression and cardiac events in non-ST-segment elevation acute coronary syndromes. *Eur Heart J*. 2005;26:2106–13.
- [96] Giffit K, Ghadban R, Assefa N, Luebbering Z, Allaham H, Enezate T. The accuracy of distribution of non-ST elevation electrocardiographic changes in localising the culprit vessel in non-ST elevation myocardial infarction. *Arch Med Sci Atheroscler Dis*. 2020;5:e226–9.
- [97] Morris N, Reynard C, Body R. The low accuracy of the non-ST-elevation myocardial infarction electrocardiograph criteria of the fourth universal definition of myocardial infarction. *Hong Kong J Emerg Med*. 2020;27:229–35.
- [98] Somers MP, Brady WJ, Perron AD, Mattu A. The prominent T wave: electrocardiographic differential diagnosis. *Am J Med*. 2002;20:243–51.
- [99] Schläpfer J, Wellens HJ. Computer-interpreted electrocardiograms: benefits and limitations. *J Am Coll Cardiol*. 2017;70:1183–92.
- [100] Wang Z, Stavrakis S, Yao B. Hierarchical deep learning with Generative Adversarial Network for automatic cardiac diagnosis from ECG signals. *Comput Biol Med*. 2023;155:106641.
- [101] Vylegzhanina AV, Kogan AE, Katrukha IA, Koshkina EV, Bereznikova AV, Filatov VL, et al. Full-size and partially truncated cardiac troponin complexes in the blood of patients with acute myocardial infarction. *Clin Chem*. 2019;65:882–92.
- [102] Katrukha IA, Katrukha AG. Myocardial injury and the release of troponins I and T in the blood of patients. *Clin Chem*. 2021;67:124–30.
- [103] Thygesen K, Alpert JS, Jaffe AS, Simoons ML, Chaitman BR, White HD, et al. Third universal definition of myocardial infarction. *J Am Coll Cardiol*. 2012;60:1581–98.
- [104] Zhelev Z, Hyde C, Youngman E, Rogers M, Fleming S, Slade T, et al. Diagnostic accuracy of single baseline measurement of Elecsys Troponin T high-sensitive assay for diagnosis of acute myocardial infarction in emergency department: systematic review and meta-analysis. *BMJ*. 2015;350:h15.
- [105] Pickering JW, Than MP, Cullen L, Aldous S, Ter Avest E, Body R, et al. Rapid rule-out of acute myocardial infarction with a single high-sensitivity cardiac troponin T measurement below the limit of detection: a collaborative meta-analysis. *Ann Intern Med*. 2017;166:715–24.
- [106] Neumann JT, Twerenbold R, Ojeda F, Sörensen NA, Chapman AR, Shah ASV, et al. Application of high-sensitivity troponin in suspected myocardial infarction. *N Engl J Med*. 2019;380:2529–40.
- [107] Sandoval Y, Smith SW, Schulz KM, Murakami MM, Love SA, Nicholson J, et al. Diagnosis of type 1 and type 2 myocardial infarction using a high-sensitivity cardiac troponin I assay with sex-specific 99th percentiles based on the third universal definition of myocardial infarction classification system. *Clin Chem*. 2015;61:657–63.
- [108] Aw T-C, Phua S-K, Tan S-P. Measurement of cardiac troponin I in serum with a new high-sensitivity assay in a large multi-ethnic Asian cohort and the impact of gender. *Clin Chim Acta*. 2013;422:26–8.
- [109] Shah AS, Griffiths M, Lee KK, McAllister DA, Hunter AL, Ferry AV, et al. High sensitivity cardiac troponin and the under-diagnosis of myocardial infarction in women: prospective cohort study. *BMJ*. 2015;350:g7873.
- [110] Boeddinghaus J, Nestelberger T, Koechlin L, Wussler D, Lopez-Ayala P, Walter JE, et al. Early diagnosis of myocardial infarction with point-of-care high-sensitivity cardiac troponin I. *J Am Coll Cardiol*. 2020;75:1111–24.
- [111] Apple FS, Smith SW, Greenslade JH, Sandoval Y, Parsonage W, Ranasinghe I, et al. Single high-sensitivity point-of-care whole-blood cardiac troponin I measurement to rule out acute myocardial infarction at low risk. *Circulation*. 2022;146:1918–29.
- [112] Pickering JW, Young JM, George PM, Watson AS, Aldous SJ, Troughton RW, et al. Validity of a novel point-of-care troponin assay for single-test rule-out of acute myocardial infarction. *JAMA Cardiol*. 2018;3:1108–12.
- [113] Sörensen NA, Neumann JT, Ojeda F, Giannitsis E, Spanuth E, Blankenberg S, et al. Diagnostic evaluation of a high-sensitivity troponin I point-of-care assay. *Clin Chem*. 2019;65:1592–601.
- [114] Januzzi JL, Mahler SA, Christenson RH, Rymer J, Newby LK, Body R, et al. Recommendations for institutions transitioning to high-sensitivity troponin testing. *J Am Coll Cardiol*. 2019;73:1059–77.
- [115] Aakre KM, Saenger AK, Body R, Collinson P, Hammarsten O, Jaffe AS, et al. Analytical considerations in deriving 99th percentile upper reference limits for high-sensitivity cardiac troponin assays: Educational recommendations from the IFCC Committee on Clinical Application of Cardiac Bio-Markers. *Clin Chem*. 2022;68:1022–30.
- [116] Lee KK, Ferry AV, Anand A, Strachan FE, Chapman AR, Kimenai DM, et al. Sex-specific thresholds of high-sensitivity troponin in patients with suspected acute coronary syndrome. *J Am Coll Cardiol*. 2019;74:2032–43.
- [117] Greene DN, Schmidt RL, Christenson RH, Rongitsch J, Imborek KL, Rebuck H, et al. Distribution of high-sensitivity cardiac troponin and N-terminal pro-brain natriuretic peptide in healthy transgender people. *JAMA Cardiol*. 2022;7:1170–4.
- [118] Welsh P, Preiss D, Hayward C, Shah ASV, McAllister D, Briggs A, et al. Cardiac troponin T and troponin I in the general population. *Circulation*. 2019;139:2754–64.
- [119] Hickman PE, Abhayaratna WP, Potter JM, Koerbin G. Age-related differences in hs-cTnI concentration in healthy adults. *Clin Biochem*. 2019;69:26–9.
- [120] Welsh P, Preiss D, Shah ASV, McAllister D, Briggs A, Boachie C, et al. Comparison between high-sensitivity cardiac troponin T and cardiac troponin I in a large general population cohort. *Clin Chem*. 2018;64:1607–16.
- [121] Collinson P, Aakre KM, Saenger A, Body R, Hammarsten O, Jaffe AS, et al. Cardiac troponin measurement at the point of care: educational recommendations on analytical and clinical aspects by the IFCC Committee on Clinical Applications of Cardiac Bio-Markers (IFCC C-CB). *Clin Chem Lab Med*. 2023;61:989–98.
- [122] Collinson PO, Saenger AK, Apple FS. High sensitivity, contemporary and point-of-care cardiac troponin assays: educational aids developed by the IFCC Committee on Clinical Application of Cardiac Bio-Markers. *Clin Chem Lab Med*. 2019;57:623–32.
- [123] Cullen L, Greenslade J, Parsonage W, Stephensen L, Smith SW, Sandoval Y, et al. Point-of-care high-sensitivity cardiac troponin in