

METHODS

The methods used to perform this systematic review were employed to minimize bias and enhance transparency in the selection, appraisal, and analysis of the available evidence. These processes are vital to the development of reliable, transparent, and accurate clinical recommendations. To view the full AAOS clinical practice guideline methodology please visit <https://www.aaos.org/additonalresources/>.

This clinical practice guideline evaluates the management of osteoarthritis of the knee (non-arthroplasty) patient outcomes. The AAOS approach incorporates practicing physicians (clinical experts) and methodologists who are free of potential conflicts of interest relevant to the topic under study, as recommended by clinical practice guideline development experts.¹

This clinical practice guideline was prepared by the AAOS Osteoarthritis of the Knee Guideline physician development group (clinical experts) with the assistance of the AAOS Clinical Quality and Value (CQV) Department (methodologists). To develop this clinical practice guideline, the clinical practice guideline development group held an introductory meeting on February 23, 2018 to establish the scope of the clinical practice guideline. As the physician experts, the clinical practice guideline development group defined the scope of the clinical practice guideline by creating PICO Questions (i.e. population, intervention, comparison, and outcome) that directed the literature search. The AAOS Medical Librarian created and executed the search (see Appendix III for search strategy).

LITERATURE SEARCHES

We begin the systematic review with a comprehensive search of the literature. Articles we consider were published prior to the start date of the search in a minimum of three electronic databases; PubMed, EMBASE, and the Cochrane Central Register of Controlled Trials. The medical librarian conducts the search using key terms determined from the guideline development group's PICO questions.

A CQV methodologist will review/include only primary literature but will supplement the electronic search with a manual search of the bibliographies of secondary literature sources, such as systematic reviews, as available. The methodologist will then evaluate all recalled articles for possible inclusion based on the study selection criteria and will summarize the evidence for the guideline work group who assist with reconciling possible errors and omissions.

A study attrition diagram is provided in the appendix of each document that details the numbers of identified abstracts, recalled and selected studies, and excluded studies that were evaluated in the CPG. The search strategies used to identify the abstracts is also included in the appendix of each CPG document.

DEFINING THE STRENGTH OF RECOMMENDATION

Judging the quality of evidence is only a steppingstone towards arriving at the strength of a CPG recommendation. The strength of recommendation also takes into account the quality, quantity, and the trade-off between the benefits and harms of a treatment, the magnitude of a treatment's effect, and whether data exists on critical outcomes.

Strength of recommendation expresses the degree of confidence one can have in a recommendation. As such, the strength expresses how possible it is that a recommendation will be overturned by future evidence. It is very difficult for future evidence to overturn a recommendation that is based on many high quality randomized controlled trials that show a large effect. It is much more likely that future evidence will overturn recommendations derived from a few small retrospective comparative studies.

Consequently, recommendations based on the former kind of evidence are given a "strong" strength of recommendation and recommendations based on the latter kind of evidence are given a "limited" strength.