

- Roquelaure,Y., Mariel,J., Dano,C., Fanello,S., Penneau-Fontbonne,D. Prevalence, incidence and risk factors of carpal tunnel syndrome in a large footwear factory. *Int.J Occup.Med Environ.Health* 2001; 4: 357-367
- Sabry,M.M., Elkader,G.A., Fahmi,M.K., Abdel-Rehman,A. Correlation of nerve conduction studies to the anthropometric measurements of the hand and to the clinical severity of Carpal Tunnel syndrome. *Egyptian Journal of Neurology, Psychiatry and Neurosurgery* 2009; 1: 67-77
- Saeed,F.-U., Hanif,S., Aasim,M. The effects of laser and ultrasound therapy on carpal tunnel syndrome. *Pakistan Journal of Medical and Health Sciences* 2012; 1: 238-241
- Saw,N.L., Jones,S., Shepstone,L., Meyer,M., Chapman,P.G., Logan,A.M. Early outcome and cost-effectiveness of endoscopic versus open carpal tunnel release: a randomized prospective trial. *J Hand Surg Br* 2003/10; 5: 444-449
- Sennwald,G.R., Benedetti,R. The value of one-portal endoscopic carpal tunnel release: a prospective randomized study. *Knee Surg Sports Traumatol.Arthrosc.* 1995; 2: 113-116
- Sharifi-Mollayousefi,A., Yazdchi-Marandi,M., Ayramlou,H., Heidari,P., Salavati,A., Zarrintan,S., Sharifi-Mollayousefi,A. Assessment of body mass index and hand anthropometric measurements as independent risk factors for carpal tunnel syndrome. *Folia Morphol.(Warsz.)* 2008/2; 1: 36-42
- Sheean,G.L., Houser,M.K., Murray,N.M. Lumbrical-interosseous latency comparison in the diagnosis of carpal tunnel syndrome. *Electroencephalogr.Clin Neurophysiol.* 1995/12; 6: 285-289
- Shin,J., Nishioka,M., Shinko,S., Shibuya,K., Sugiki,M., Kasumoto,H., Fudo,A., Bito,Y., Fujita,Y., Komaba,K. Carpal tunnel syndrome and plasma beta2-microglobulin concentration in hemodialysis patients. *Ther Apher.Dial.* 2008/2; 1: 62-66
- Shiota,E., Tsuchiya,K., Yamaoka,K., Kawano,O. Open surgical therapy for carpal tunnel decompression in long-term haemodialysis patients. *Journal of Hand Surgery* 2001; 6: 529-532
- Shum,C., Parisien,M., Strauch,R.J., Rosenwasser,M.P. The role of flexor tenosynovectomy in the operative treatment of carpal tunnel syndrome. *J Bone Joint Surg Am* 2002/2; 2: 221-225
- Silverstein,B.A., Fine,L.J., Armstrong,T.J. Occupational factors and carpal tunnel syndrome. *Am J Ind.Med* 1987; 3: 343-358
- Smith,T. Near-nerve versus surface electrode recordings of sensory nerve conduction in patients with carpal tunnel syndrome. *Acta Neurol Scand.* 1998/10; 4: 280-282
- Sorensen,A.M., Dalsgaard,J., Hansen,T.B. Local anaesthesia versus intravenous regional anaesthesia in endoscopic carpal tunnel release: a randomized controlled trial. *J Hand Surg Eur.Vol.* 2013/6; 5: 481-484
- Soyupek,F., Yesildag,A., Kutluhan,S., Askin,A., Ozden,A., Uslusoy,G.A., Demirci,S. Determining the effectiveness of various treatment modalities in carpal tunnel syndrome by ultrasonography and comparing ultrasonographic findings with other outcomes. *Rheumatol.Int.* 2012/10; 10: 3229-3234
- Stalberg,E., Stalberg,S., Karlsson,L. Automatic carpal tunnel syndrome tester. *Clin Neurophysiol.* 2000/5; 5: 826-832