

MRI can image fascia, vessels, nerve, muscle, cartilage, ligaments, tendons, pannus, synovial effusions, and bone marrow. Visualization of particular structures can be enhanced by altering the pulse sequence to produce either T1- or T2-weighted spin echo, gradient echo, or inversion recovery (including short tau inversion recovery [STIR]) images. Because of its sensitivity to changes in marrow fat, MRI is a sensitive but nonspecific means of detecting osteonecrosis, osteomyelitis, and marrow inflammation indicating overlying synovitis or osteitis (Fig. 363-8). Because of its enhanced soft tissue resolution, MRI is more sensitive than arthrography or CT in the diagnosis of soft tissue injuries (e.g., meniscal and rotator cuff tears); intraarticular derangements; marrow abnormalities (osteonecrosis, myeloma); and spinal cord or nerve root damage, synovitis, or cartilage damage or loss.

ACKNOWLEDGMENT

The author acknowledges the insightful contributions of Dr. Peter E. Lipsky to this chapter in previous editions.

■ FURTHER READING

ALI Y: Rheumatologic tests: A primer for family physicians. *Am Fam Physician* 98:164, 2018.

CUSH JJ et al: Evaluation of musculoskeletal complaints. Available from <http://www.rheumaknowledgy.com/evaluation-of-musculoskeletal-complaints>. Accessed April 6, 2017.

HUBBARD MJ et al: Common soft tissue musculoskeletal pain disorders. *Prim Care* 45:289, 2018.

OLSEN NJ, KARP DR: Finding lupus in the ANA haystack. *Lupus Sci Med* 7:e000384, 2020.

RUDWALEIT M et al: How to diagnose axial spondyloarthritis early. *Ann Rheum Dis* 63:535, 2004.

SIMPFENDORFER CS: Radiologic approach to musculoskeletal infections. *Infect Dis Clin North Am* 31:299, 2017.